



U.S. Fish and Wildlife Service

# Distribution and habitat use of fish in Seattle's streams

*Final Report, 2005 and 2006*

*March 2009*

*By Daniel W. Lantz, Roger A. Tabor, and Scott T. Sanders*

*U.S. Fish and Wildlife Service  
Washington Fish & Wildlife Office  
Lacey, Washington*



Seattle  
Public  
Utilities

*Funded by Seattle Public Utilities (City of Seattle)*



**Distribution and habitat use of fish in Seattle's streams  
Final Report, 2005 and 2006**

by

Daniel W. Lantz, Roger A. Tabor, and Scott T. Sanders.

U.S. Fish and Wildlife Service  
Washington Fish and Wildlife Office  
Fisheries Division  
510 Desmond Drive SE, Suite 102  
Lacey, Washington 98503

March 2009



## EXECUTIVE SUMMARY

In 2005 and 2006, we conducted a comprehensive survey of Seattle's streams to determine fish distribution. Information from this inventory will aid the City of Seattle with their future management decisions. The survey consisted of two major parts: 1) a survey of all streams to determine the overall distribution of each fish species (single-pass electrofishing), and 2) surveys of long-term reference sites (multiple-pass electrofishing) to estimate fish abundance. Fish distribution surveys were conducted during the summer and winter. Reference site surveys were conducted in at least one site of the five major watersheds (Piper's Creek, Thornton Creek, Longfellow Creek, Fauntleroy Creek, and Taylor Creek). Fish communities in the reference sites were sampled for abundance, biomass, and diversity.

During both parts of the study, we also collected stream habitat information. Stream habitat surveys were conducted on 37 out of the 49 stream systems. Of the 37 streams surveyed, we collected habitat data on a total of 149 sites. In general, habitat conditions appeared good in Piper's Creek, Thornton Creek, and Longfellow Creek. In Taylor Creek and Fauntleroy Creek, many sections were shallow and had little pool habitat. In the smaller stream systems, habitat conditions often did not appear conducive to supporting fish populations. Often much of the stream was in a long culvert and may be a barrier to upstream fish movements. The wetted stream width was often narrow and there was little streamflow. These streams tended to be shallow (maximum depth < 0.25 m) and have little pool habitat. Also, the substrate was predominantly sand with little gravel or larger substrates.

Summer surveys, indicated cutthroat trout *Oncorhynchus clarkii* were widespread in Piper's Creek, Thornton Creek, and Taylor Creek. Only one cutthroat trout was ever collected in Longfellow Creek despite a large amount of available habitat. Additionally, cutthroat trout were absent in other southwest stream systems. Additional research is needed to better understand why cutthroat trout are rare in the southwest streams including Longfellow Creek but abundant in other similar-sized streams. Winter surveys of Piper's Creek and South Branch of Thornton Creek documented the presence of cutthroat trout in more upstream locations.

Juvenile coho salmon *O. kisutch* were observed in all of the five major watersheds as well as Durham Creek; however, it's unclear if they were naturally-produced or were part of an enhancement project. Rainbow trout *O. mykiss* were rarely collected and were only observed in Thornton Creek, Ravenna Creek, Longfellow Creek, and Puget Creek. The only location we ever observed juvenile Chinook salmon *O. tshawytscha* was in Taylor Creek, near its mouth on Lake Washington.

Threespine stickleback *Gasterosteus aculeatus* were also widespread in Seattle stream systems but were usually found in large numbers in ponds or low-velocity areas of streams. The only freshwater species of cottids found in Seattle streams were coastrange sculpin *Cottus aleuticus* and prickly sculpin *C. asper*. Cottids were only found in the low reaches of streams. Because of their poor swimming ability and they undergo a pelagic

larvae phase in downstream areas (Lake Washington or Puget Sound), their ability to inhabit upstream areas can be limited by small instream barriers such as small cascades and weirs. Introduced fish species observed included four centrarchid species and four other species. Introduced species were primarily observed in the Thornton Creek mainstem and the North Branch of Thornton Creek.

A total of nine reference sites were established in the major watersheds. Salmonid biomass estimates in Thornton Creek and Piper's Creek drainages were generally higher than the other three systems. The density of salmonids in our reference sites in the South Branch of Thornton Creek and Piper's Creek appeared to be high in comparison to other lowland streams in the Pacific Northwest.

To assess ecosystem health, we used a fish index of biotic integrity (FIBI) that has been developed for other Puget Sound lowland streams. FIBI scores were generally low in Piper's Creek and Thornton Creek watersheds, primarily due to the relatively high abundance of cutthroat trout and lack of other species such as coho salmon and cottids. FIBI scores were generally higher in Longfellow Creek and Fauntleroy Creek than other Seattle streams, largely because few cutthroat trout were present and coho salmon made up a high percentage of the catch. However, the FIBI scores from these streams may have been artificially high because juvenile coho salmon may have been outplanted.

## TABLE OF CONTENTS

|   |     |
|---|-----|
| INTRODUCTION .....                              | 1   |
| CHAPTER 1: HABITAT.....                         | 6   |
| Introduction and Methods .....                  | 6   |
| Results.....                                    | 7   |
| CHAPTER 2: DISTRIBUTING AND TYPING.....         | 29  |
| Introduction and Methods .....                  | 29  |
| Fish Distribution Surveys .....                 | 29  |
| Stream Typing.....                              | 30  |
| Results.....                                    | 34  |
| Fish Distribution by Region.....                | 34  |
| Fish Distribution by Species .....              | 45  |
| Stream Typing.....                              | 60  |
| Discussion.....                                 | 62  |
| CHAPTER 3: REFERENCE SITE FISH DENSITIES .....  | 74  |
| Introduction and Methods .....                  | 74  |
| Results.....                                    | 76  |
| Discussion.....                                 | 89  |
| CHAPTER 4: FISH INDEX OF BIOTIC INTEGRITY ..... | 92  |
| Introduction and Methods.....                   | 92  |
| Results.....                                    | 93  |
| Discussion.....                                 | 95  |
| ACKNOWLEDGEMENTS.....                           | 97  |
| REFERENCES .....                                | 98  |
| APPENDIX.....                                   | 103 |





## INTRODUCTION

The presence or absence of fishes in streams is an important determinant for stream and riparian zone management regimes. In Washington State, streams are often classified according to the presence of fishes, particularly anadromous salmonids. Resource management decisions are guided by these resulting stream classifications. The stream classification system is used in urban streams as well as rural and forested stream environments. In 2005 and 2006, we conducted a comprehensive survey of Seattle's streams to determine the presence, abundance, and diversity of fish. The survey consisted of two major parts: a survey of all streams to determine the overall distribution of each fish species (single-pass electrofishing), and surveys of long-term reference sites (multiple-pass electrofishing) to estimate fish abundance. The first fish distribution surveys were conducted from June to October 2005. Additional surveys were conducted in February 2006 to determine if fish distributions changed according to season. Reference site surveys were conducted in the winter and fall in 2005 and during the summer in 2006. Surveys were conducted in the five major watersheds. Fish communities in the reference sites were sampled for abundance, biomass, and diversity.

Seattle Public Utilities has identified 49 urban watersheds within the Seattle City limits (Figure 1). Of the 49 watersheds, five (Piper's Creek, Thornton Creek, Longfellow Creek, Fauntleroy Creek, and Taylor Creek) are considered major watersheds based on the size of the watershed and amount of available stream habitat. In 1999, Wild Fish Conservancy (Washington Trout 2000) conducted an initial stream typing and fish barrier survey of these five creeks and other streams. However, surveys were only conducted during the summer period and most of the small creeks were not surveyed. In addition, some fishes, such as sculpins (Table 1), were not identified to species and the distribution of some species may not have been underestimated. Prior to the Washington Trout (2000) survey, most electrofishing surveys were conducted in either Thornton Creek or Piper's Creek. Results of surveys from Thornton Creek (Muto and Shefler 1983; Ludwa et al. 1997) indicated cutthroat trout were abundant and few other fish were present. Surveys of Piper's Creek by Pfeifer (1984) and Thomas (1992) found cutthroat trout,

## INTRODUCTION

The presence or absence of fishes in streams is an important determinant for stream and riparian zone management regimes. In Washington State, streams are often classified according to the presence of fishes, particularly anadromous salmonids. Resource management decisions are guided by these resulting stream classifications. The stream classification system is used in urban streams as well as rural and forested stream environments. In 2005 and 2006, we conducted a comprehensive survey of Seattle's streams to determine the presence, abundance, and diversity of fish. The survey consisted of two major parts: a survey of all streams to determine the overall distribution of each fish species (single-pass electrofishing), and surveys of long-term reference sites (multiple-pass electrofishing) to estimate fish abundance. The first fish distribution surveys were conducted from June to October 2005. Additional surveys were conducted in February 2006 to determine if fish distributions changed according to season. Reference site surveys were conducted in the winter and fall in 2005 and during the summer in 2006. Surveys were conducted in the five major watersheds. Fish communities in the reference sites were sampled for abundance, biomass, and diversity.

Seattle Public Utilities has identified 49 urban watersheds within the Seattle City limits (Figure 1). Of the 49 watersheds, five (Piper's Creek, Thornton Creek, Longfellow Creek, Fauntleroy Creek, and Taylor Creek) are considered major watersheds based on the size of the watershed and amount of available stream habitat. In 1999, Wild Fish Conservancy (Washington Trout 2000) conducted an initial stream typing and fish barrier survey of these five creeks and other streams. However, surveys were only conducted during the summer period and most of the small creeks were not surveyed. In addition, some fishes, such as sculpins (Table 1), were not identified to species and the distribution of some species may not have been underestimated. Prior to the Washington Trout (2000) survey, most electrofishing surveys were conducted in either Thornton Creek or Piper's Creek. Results of surveys from Thornton Creek (Muto and Shefler 1983; Ludwa et al. 1997) indicated cutthroat trout were abundant and few other fish were present. Surveys of Piper's Creek by Pfeifer (1984) and Thomas (1992) found cutthroat trout,

sculpin, and juvenile coho salmon were common in the lower reach (river kilometer [Rkm] 0-0.65), but only cutthroat trout were present in the upper reach (Rkm 6.5-1.75). In addition to fish surveys, the city has collected a wide variety of other information on their major streams. This includes: culvert assessments, habitat assessments, channel condition assessments, riparian assessments, streamflow monitoring, salmonid spawning surveys, smolt trapping, and benthic invertebrate sampling (City of Seattle 2007). Most of the monitoring and research efforts on Seattle's streams have been focused on the larger streams of the five major watersheds. Streams in the smaller watersheds and smaller tributaries of the five watersheds have received little attention. One important objective of this study was to provide the City of Seattle with habitat and fish distribution on all streams, not just the large streams of the five major watersheds.

The native ichthyofauna of Pacific Northwest streams consists largely of members of the families Salmonidae (salmon, trout, charr, and whitefish) and Cottidae (sculpins). Information on the distribution and habitat requirements of salmonids has been studied extensively. In contrast, the distribution and habitat requirements of cottids are not well known (Tabor et al. 2007). Because cottids are not economically important and they can be difficult to identify, researchers usually collect minimal information on these fish. As part of our assessment, we were interested in collecting detailed information of cottid distribution. Because they are more commonly present near the stream's mouth, we conducted supplemental surveys at these areas to accurately determine the distribution of each cottid species and identify potential barriers that may limit their upstream movement.

The streams of Seattle lie within a heavily urbanized area. Of the five major watershed, the percent of impervious surfaces ranges from 38% in the Fauntleroy Creek watershed to 59% for the Thornton Creek watershed (City of Seattle 2007). Effects of urbanization to the health of the stream ecosystem include: increased peak streamflow, reduced substrate size, reduction in large woody debris, reduction in the pool depth, increased water temperatures, increased levels of chemical contaminants, reduction in benthic invertebrate diversity and abundance, and changes in fish assemblage including

introductions of exotic species (Karr 1998). Changes to the fish assemblage often include an increase in the proportion of cutthroat trout and reduction in abundance of other salmonids and sculpin (Serl 1999).

The overall objectives of this study included:

1. Assess habitat quality to previously unsurveyed smaller streams to better inform the City's future management decisions – Chapter 1.
2. Classify Seattle's urban streams and stream reaches according to the current Washington State stream classification system – Chapter 2.
3. Determine the presence/absence, distribution, species composition, and relative abundance of fish (salmonids and non-salmonids) in all City of Seattle streams – Chapter 2.
4. Quantify fish abundance and biomass and collect habitat information at reference sites (including potential restoration sites in Thornton Creek) in the five major watersheds – Chapter 3.
5. Use a fish index of biotic integrity (FIBI) to assess ecosystem health – Chapter 4.




FIGURE 1.—Map of the City of Seattle displaying the 49 stream systems within the city limits (purple lines).

TABLE 1.-- Scientific and common names of native and nonnative fishes of Seattle's streams and ponds mentioned in this report.

| <b>Family</b><br>Genus and Species | <b>Common Name</b>           | <b>Native/Nonnative</b> |
|------------------------------------|------------------------------|-------------------------|
| <b>Petromyzontidae</b>             | Lamprey<br>(Unknown species) | Native                  |
| <b>Salmonidae</b>                  |                              |                         |
| <i>Oncorhynchus tshawytscha</i>    | Chinook salmon               | Native                  |
| <i>Oncorhynchus kisutch</i>        | Coho salmon                  | Native                  |
| <i>Oncorhynchus clarkii</i>        | Cutthroat trout              | Native                  |
| <i>Oncorhynchus mykiss</i>         | Rainbow trout                | Native                  |
| <b>Cyprinidae</b>                  |                              |                         |
| <i>Carassius auratus</i>           | Goldfish                     | Nonnative               |
| <i>Cyprinus carpio</i>             | Common carp (Koi)            | Nonnative               |
| <i>Mylocheilus caurinus</i>        | Peamouth                     | Native                  |
| <b>Cobitidae</b>                   |                              |                         |
| <i>Misgurnus anguillicaudatus</i>  | Oriental weatherfish         | Nonnative               |
| <b>Ictaluridae</b>                 |                              |                         |
| <i>Ameriurus nebulosus</i>         | Brown bullhead               | Nonnative               |
| <b>Gasterosteidae</b>              |                              |                         |
| <i>Gasterosteus aculeatus</i>      | Threespine stickleback       | Native                  |
| <b>Centrarchidae</b>               |                              |                         |
| <i>Micropterus salmonides</i>      | Largemouth bass              | Nonnative               |
| <i>Micropterus dolomieu</i>        | Smallmouth bass              | Nonnative               |
| <i>Ambloplites rupestris</i>       | Rock bass                    | Nonnative               |
| <i>Lepomis gibbosus</i>            | Pumpkinseed                  | Nonnative               |
| <b>Cottidae</b>                    |                              |                         |
| <i>Cottus aleuticus</i>            | Coastrange sculpin           | Native                  |
| <i>Cottus asper</i>                | Prickly sculpin              | Native                  |
| <i>Clinocottus acuticeps</i>       | Sharpnose sculpin            | Native                  |
| <i>Leptocottus armatus</i>         | Pacific staghorn sculpin     | Native                  |
| <b>Ammodytidae</b>                 |                              |                         |
| <i>Ammodytes hexapterus</i>        | Pacific sand lance           | Native                  |

## CHAPTER 1. STREAM PHYSICAL HABITAT SURVEYS

### Introduction and Methods

Past habitat surveys of Seattle's streams have focused on the large streams of the five major watersheds. Little attention has been given to smaller streams. Information on these other streams will allow managers to make informed decisions on land-use management and potential restoration projects. As part of our fish distribution surveys, we collected stream habitat information at all the sites we surveyed, which included both small streams and large streams of the five major watersheds.

Stream habitat surveys were conducted on most streams during the fish distribution surveys (see Chapter 2). We attempted to survey all streams of the 49 watersheds; however, lack of access, private property issues, or general inaccessibility prevented us from surveying a few of the smaller streams. Streams that were accessible were first surveyed for fish use and then physical habitat. For convenience, the 49 watersheds were grouped into six geographical areas (Figure 1). At each site, we divided the stream into habitat units, generally following the procedures of Pleus et al. (1999), except we classified some habitat units as glides, which were shallow habitats with low water velocities (no turbulence). All stream habitat data measurements were measured to the nearest tenth of a meter using a hand held stadia rod or a laser rangefinder. Habitat data collected were as follows:

Length- The length of each site was measured along the thalweg from the downstream end to the upstream end of the site. Any habitat units within the site were also measured in this manner.

Wetted width- The wetted width was measured at three equidistance locations along the measured length of each habitat unit. Working upstream, a measurement was taken at  $\frac{1}{4}$  distance from the downstream end, at  $\frac{1}{2}$  of the length of the site, and at  $\frac{3}{4}$  the length of the site. These were then averaged to obtain an average wetted width.

Depth- An average and maximum depth was recorded at each habitat and each site. The average depth was measured by randomly placing the hand held stadia rod at different locations within the habitat and averaging these values. The maximum depth of the habitat was collected by simply using the stadia rod to find the deepest part of the habitat. If the habitat unit was a pool, the depth was measured at the deepest part and at deepest part of the tailout (pool outlet). By subtracting the tailout depth from the maximum pool depth, we calculated a residual pool depth. If more than one habitat type was in a single site, the maximum depth, or the deepest habitat of all the habitats was used as the maximum depth for the entire site. The average depth of a site containing multiple habitats was obtained by averaging the average depths of all the habitats.

Substrate- Substrate values were collected by visually identifying the types of substrates and then estimating the percent for each within each habitat type. Substrate values were recorded as: sand/silt, gravel, cobble, boulders, or rip rap.

Discharge or streamflow- If possible, streamflow (measured in cubic feet per second, cfs) was measured at the mouth of each stream. For medium- and large-sized streams we generally followed the procedures of Pleus (1999). We placed a metric tape across the mouth of the stream and divided the stream width into 12-20 cells. In the middle of each cell, the water depth and average water velocity was measured. Average water velocity was measured at 60% of the total depth from the surface with a Marsh-McBirney® model-2000 portable flowmeter. Stream discharge could then be calculated by the equation;

$$Q = \sum A_i V_i ;$$

where Q is discharge or streamflow;  $A_i$  is the cross sectional area [width x depth] for cell i; and  $V_i$  is the average velocity of cell i.

If the stream had little streamflow and was too shallow for the flowmeter to work, we often used a volumetric measurement method (Rantz et al. 1982) to collect the streamflow data. We tried to find a location where the stream dropped over a short waterfall (i.e., outflow of a perched culvert) and the entire flow could be captured in a bucket. At some sites, we created a small dam in the stream using sandbags and placed a 10-cm-diameter pipe on the top of the dam for the water to flow through. A bucket was quickly placed below the pipe or waterfall and was allowed to fill. After the bucket was close to being full, it was removed quickly and the amount of water in the bucket was measured. A stopwatch was used to time the duration water was flowing into the bucket. This was done at least three times to get an average streamflow.

Temperature and conductivity- We also recorded the temperature (°C), and the conductivity ( $\mu$ S, micro-Siemens) by the use of a Hanna® portable waterproof multi-range conductivity /TDS meter model HI 9635.

## **Results**

Stream habitat surveys were conducted on 37 out of the 49 stream basins. Streams #'s 2, 3, 6, 7, 11, 16, and 23-25 were not surveyed because they were inaccessible, and Streams #'s 36, 41 and 43 were not surveyed because these streams were completely dry. Of the 37 streams surveyed, we collected habitat data on a total of 149 fish distribution sites. Due to the City's already extensive habitat data on the mainstems of Piper's, Thornton, Longfellow, Fauntleroy, and Taylor Creeks, habitat data presented is focused mainly on the smaller streams throughout the city and the lesser known tributaries of the five major creeks.

*Northwest region.*-- The northwest region contains approximately 12 streams (streams system #'s 1–13) from the city's northern border with the City of Shoreline, south to and including Golden Gardens Park, and east to Licton Springs Park (Figure 1). The largest stream



in this region is Piper's Creek, which includes Mohlendorph Creek and Venema Creek and several smaller tributaries. Habitat data was collected on 3 sites on stream system #1, one of which was a 0.093 ha man-made pond (Table 1.1). Substrate in the pond was 100% sand/silt. Upstream of the pond, the stream was small, shallow, and had mostly sand/silt substrate. Downstream of the pond we surveyed a 75-m long site which was heavily landscaped and flowed through several sections with culverts. This section was high gradient with two vertical drops, one of 1.5 m and another of 7 m. The substrate in this section appeared to be all cobble.

TABLE 1.1.-- Size and maximum depth of ponds sampled in Seattle, 2005-2006. Appendix number is the map which displays the location of each site. For irregular shaped ponds, more than one width is given.

| Stream system #            | Region<br>Pond name             | Appendix | Area (ha) | Length (m) | Widths (m) | Maximum depth (m) |
|----------------------------|---------------------------------|----------|-----------|------------|------------|-------------------|
| <b>Northwest region</b>    |                                 |          |           |            |            |                   |
| 1                          | Unnamed stream # 1 pond         | 1.1      | 0.093     | 27.4       | 34.1       | 4.0               |
| <b>Northeast region</b>    |                                 |          |           |            |            |                   |
| 14                         | Matthews Creek pond             | 1.8      | 0.035     | 22         | 16         | 1.5               |
| 14                         | Littles Creek                   | 1.13     | 0.364     | 192.5      | 25         | --                |
| 14                         | Jackson Park Golf Course A pond | 1.13     | 0.445     | 107        | 55         | 2.1               |
| 14                         | Jackson Park Golf Course B pond | 1.13     | 0.342     | 113        | 49         | 2.7               |
| 14                         | Jackson Park Golf Course C pond | 1.13     | 0.586     | 175        | 19,50      | --                |
| 14                         | Park and Ride pond              | 1.16     | 0.024     | 24         | 10         | 1.0               |
| 14                         | NSCC pond                       | 1.16     | 0.437     | 196        | 31         | --                |
| <b>West-central region</b> |                                 |          |           |            |            |                   |
| 21                         | Scheuerman Creek pond           | 1.20     | 0.030     | 24.5       | 12.3       | 1.3               |
| 21                         | Scheuerman Creek trib pond A    | 1.20     | 0.035     | 22         | 16.1       | 1.5               |
| 21                         | Scheuerman Creek trib pond B    | 1.20     | 0.033     | 25.4       | 12.9       | 1.0               |
| 21                         | Scheuerman Creek trib pond C    | 1.20     | 0.021     | 21         | 10         | 2.0               |
| 22                         | Sewer Plant Beach Pond          | 1.20     | 0.219     | 220        | 8,21,11    | 1.5               |
| 22                         | Unnamed Trading Post Pond       | 1.20     | 0.074     | 30.6       | 24.2       | 0.75              |
| <b>East-central region</b> |                                 |          |           |            |            |                   |
| 27                         | Japanese Gardens pond           | 1.21     | 0.198     | 95         | 13,25,25   | 1.5               |
| 27                         | Unnamed trib pond               | 1.21     | 0.070     | 26         | 27         | 1.2               |
| <b>Southeast region</b>    |                                 |          |           |            |            |                   |
| 48                         | Kabota Gardens pond A           | 1.29     | 0.012     | 15         | 8          | --                |
| 48                         | Kabota Gardens pond B           | 1.29     | 0.040     | 50         | 8          | --                |
| 48                         | Kabota Gardens pond C           | 1.29     | 0.014     | 17         | 8          | --                |

Habitat data was collected from a 3.1-m long section of stream system #4 that ran along a private drive near the stream mouth. The average depth at this site was 0.01 m with a maximum depth of 0.02 m. Substrate was 15 % sand/silt and 85 % gravel.

On Piper's Creek (stream system #5), we collected habitat data from four sites from three small unnamed tributaries. On the mainstem of Piper's Creek, we surveyed a combined total length of 510 m, with an average width of 2.6 m, an average depth of 0.1 m and an average maximum depth of 0.3 m, with the deepest recorded depth of 0.75 m (Table 1.2). Riffles were the dominant habitat type comprising 50%, along with glides 24 %, and pools 26 %. Substrate in Piper's Creek was composed mostly of fines and gravel (Table 1.3). Several smaller tributaries of Piper's Creek were also surveyed. The tributaries adjacent to the treatment plant and immediately downstream were just small springs emanating from the hillside (Figure 1.1) and had no fish habitat. Upstream of the treatment plant, we surveyed three small tributaries, each of which had a high percentage of gravel and cobble (Table 1.3).

Or the most part, Mohlendorph Creek was dry during the summer 2005 survey and was resurveyed in the winter of 2006. A 12.1 m long section starting at the mouth of Mohlendorph Creek was surveyed which had an average width of 1.6 m, an average depth of 0.09 m with a max depth of 0.25 m, which was taken from the pool at the confluence with Venema Creek (Figure 1.2). Substrate in Mohlendorph Creek, upstream of the weir, was primarily composed of gravel with some sand and cobble. Venema Creek was surveyed 600 m upstream from the confluence with Piper's Creek. At this site we sampled a 50 m section comprised of two riffles and one pool, which was the deepest part of this site. Substrate was composed mostly of sand with some gravel (Table 1.3).

TABLE 1.2.-- Stream habitat data for streams surveyed in the northwest region, October 2005. Locations of sample sites are displayed in Appendices 1.1-1.7. Max = maximum. Percent riffles and pools/glides are the percent of the total stream area.

| Stream system # | Sample # | Stream Name             | Date Surveyed | Distance from mouth (m) | Mean wetted width (m) | Mean depth (m) | Max depth (m) | % riffles | % pools/glides |
|-----------------|----------|-------------------------|---------------|-------------------------|-----------------------|----------------|---------------|-----------|----------------|
| 1               | 3        | Unnamed                 | 4-Oct         | 170-216                 | 1.15                  | 0.05           | 0.11          | 100.0     | 0.0            |
| 4               | 4        | Broadview               | 4-Oct         | 205-208                 | 0.36                  | 0.01           | 0.02          | 100.0     | 0.0            |
| 5               | 25       | Piper's Creek           | 6-Oct         | 0-42                    | 3.20                  | 0.15           | 0.4           | 0.0       | 100.0          |
| 5               | 6        | Piper's Creek           | 6-Oct         | 42-74                   | 2.50                  | 0.07           | 0.2           | 23.1      | 76.9           |
| 5               | 7        | Piper's Creek           | 6-Oct         | 250-300                 | 3.05                  | 0.11           | 0.75          | 55.2      | 44.8           |
| 5               | 8        | Piper's Creek           | 6-Oct         | 550-600                 | 3.89                  | 0.21           | 0.75          | 73.8      | 26.2           |
| 5               | 18       | Venema Creek            | 20-Oct        | 600-650                 | 1.07                  | 0.07           | 0.4           | 92.2      | 7.8            |
| 5               | 21       | Unnamed trib H          | 11-Oct        | 0-50                    | 1.13                  | 0.08           | 0.22          | 70.3      | 29.7           |
| 5               | 22       | Unnamed trib H          | 11-Oct        | 250-300                 | 1.32                  | 0.10           | 0.4           | 37.1      | 62.9           |
| 5               | 10       | Piper's Creek           | 6-Oct         | 1,300-1,353             | 2.65                  | 0.18           | 0.54          | 45.9      | 54.1           |
| 5               | 23       | Unnamed trib K          | 11-Oct        | 0-54                    | 1.00                  | 0.05           | 0.14          | 100.0     | 0.0            |
| 5               | 11       | Piper's Creek           | 11-Oct        | 1,600-1,650             | 2.61                  | 0.14           | 0.26          | 73.8      | 26.2           |
| 5               | 12       | Piper's Creek           | 11-Oct        | 1,750-1,800             | 1.60                  | 0.08           | 0.24          | 47.5      | 52.5           |
| 5               | 24       | Unnamed trib M          | 11-Oct        | 0-7.5                   | 0.50                  | 0.04           | 0.06          | 100.0     | 0.0            |
| 5               | 13       | Piper's Creek           | 11-Oct        | 2,150-2,181             | 0.80                  | 0.02           | 0.1           | 0.0       | 100.0          |
| 8               | 33       | Blue Ridge Creek        | 25-Oct        | 1,050-1,060             | 1.00                  | 0.03           | 0.1           | 100.0     | 0.0            |
| 8               | 30       | East Fork Blue Ridge Cr | 25-Oct        | 50-70                   | 0.50                  | 0.05           | 0.19          | 100.0     | 0.0            |
| 12              | 42       | Golden Gardens Stream E | 14-Oct        | 20-81                   | 1.25                  | 0.02           | 0.05          | 100.0     | 0.0            |
| 13              | 45       | Licton Springs          | 25-Oct        | 1,600-1,622             | 2.23                  | 0.11           | 1.1           | 10.4      | 89.6           |

TABLE 1.3.-- Total combined substrate scores for streams in the northwest region.

| Stream system # | Stream name             | # of sites | Overall Substrate Score % |        |        |          |         |
|-----------------|-------------------------|------------|---------------------------|--------|--------|----------|---------|
|                 |                         |            | Silt/Sand                 | Gravel | Cobble | Boulders | Rip Rap |
| 1               | Unnamed                 | 3          | 58                        | 2      | 40     |          |         |
| 4               | Broadview Creek         | 1          | 15                        | 85     |        |          |         |
| 5               | Piper's Creek           | 9          | 42                        | 39     | 8      | 11       |         |
| 5               | Venema Creek            | 1          | 60                        | 23     | 15     | 2        |         |
| 5               | Mohlendorph Creek       | 1          | 15                        | 70     | 15     |          |         |
| 5               | Unnamed trib H          | 2          | 26                        | 39     | 26     | 3        | 6       |
| 5               | Unnamed trib K          | 1          | 50                        | 50     |        |          |         |
| 5               | Unnamed trib M          | 1          | 5                         | 60     | 35     |          |         |
| 8               | Blue Ridge Creek        | 2          | 80                        | 20     |        |          |         |
| 9               | Unnamed                 | 1          | 50                        | 50     |        |          |         |
| 10              | Unnamed                 | 1          | 100                       |        |        |          |         |
| 12              | Golden Gardens Stream B | 2          | 50                        |        |        |          | 50      |
| 12              | Golden Gardens Stream E | 1          | 100                       |        |        |          |         |
| 13              | Licton Springs Creek    | 2          | 87                        | 12     |        |          | 1       |



FIGURE 1.1.-- Photo taken of unnamed tributary F of Piper's Creek entering downstream of the treatment plant.



FIGURE 1.2.-- Photo taken in the winter of 2006 looking upstream at the weir pool on Mohlendorph Creek. This was the deepest spot recorded on Mohlendorph Creek. This location was dry during the summer sampling of 2005. The stream flows from the upper-center of the photo, through the weir, and into the pool at the bottom of the photo.

Stream system #'s 8-10 were generally shallow ( $< 0.1$  m average depth) and were dominated primarily by fine sediment (Table 1.3; Figure 1.3). Stream systems #'s 8-10 also had large amounts of introduced plant species throughout their basins. Stream system #12 (A-G) also had high amounts of fine sediment and was overall very shallow (Table 1.3). Habitat surveys of Licton Spring, (stream system #13), was taken at two sites within Licton Springs Park. The site located at rkm 1.6 was 22.5 m long with an average width of 2.4 m, an average depth of 0.3 m with a max depth of 1.1 m, which was recorded at culvert at the south end of the park (Figure 1.4). We also surveyed upstream of this point at rkm 1.8 in a much deeper (maximum depth, 0.9 m) and wider (wetted width, 2.0 m) section.



FIGURE 1.3.-- Photos of stream system # 8. The left photo was taken in Homewood Park showing the overall shallow depth of the stream at this site and dominant fine substrate. The right photo is of the East Fork tributary of stream system # 8. The stream at this point is down cut more than 0.5 m.



FIGURE 1.4.-- Photo taken of the culvert that Licton Springs Creek flows into at the south end of Licton Springs Park. Note there are two branches flowing into the culvert. At the time of our survey, water was only flowing in from the top branch in the photo.

*Northeast region.*-- The northeast region consists primarily of the streams in the Thornton Creek drainage (stream system #14). In this drainage, we surveyed five sites on the mainstem, five sites on the North Branch, and six sites on the South Branch. Within the Thornton Creek watershed we were able to survey 35 sites on 18 separate tributaries.

The mainstem of Thornton Creek (Rm 0-2,300) is a low gradient reach with primarily sand or gravel substrate (Table 1.4). In some locations, the banks were armored with rip rap and the boulders of the rip rap were an important component of the overall substrate composition. The mainstem has a good mixture of riffles and large, deep pools (Table 1.5).

Close to the mouth of Thornton Creek, there are several small tributaries that converge and enter the main stem of Thornton Creek downstream of Sand Point Way. Habitat in this area has largely been influenced by human development and many of these small tributaries flow through several culverts before entering Thornton Creek. Matthews Creek, the lowest downstream tributary of Thornton Creek, flows approximately 120 m from a culvert and through a pond before it enters Thornton Creek. We surveyed the pond close to the mouth of Matthews Creek which was 16-m long by 22-m wide and 1.5 m at its deepest point. Upstream of the pond we surveyed a small pool and glide at Rm 120 (Figure 1.5). The pool was 6.1-m long by 7.5-m wide with a maximum depth of 0.5 m. The glide was 7.2-m long by 1.2-m wide with a maximum depth of 0.32 m and surrounded on both sides by rip rap. Substrate was composed of 80% fines, 10% gravel, and 10% rip rap for both habitats. Further upstream, where Matthews Creek enters the culvert, at the Sand Point Country Club Golf Course, the stream was dry.

On Thornton Creek about 300 m upstream from the mouth at Lake Washington is the confluence of Maple Creek. We surveyed two sites on the mainstem of Maple Creek, one at the mouth and a second 400 m upstream from the mouth on the upstream of the culvert that is under 45<sup>th</sup> Ave NE. At the mouth of Maple Creek, there is a small cascade of about 0.75 m in height. Upstream of this point the creek flows from a culvert under Sand Point Way NE and forms a small pool/glide that is approximately 2 m wide and 5 m long and dominated by fine substrate (Figure 1.6). Maple Creek then becomes a small (< 1.0 m wetted width), shallow (0.1 m, maximum depth) riffle and flows through private property before entering Thornton Creek. At the upper site, we sampled a 50-m long riffle upstream of the culvert under 45<sup>th</sup> Ave NE that was 1.0-m wide with a maximum depth of 0.13 m. Substrate composition was 50% sand and 50% gravel. A culvert at the downstream end was perched 1.2 m above the creek.

TABLE 1.4.-- Total combined substrate scores for streams in the northeast region.

| Stream system # | Stream name                 | # of Sites | Overall Substrate Score % |        |        |          |         |
|-----------------|-----------------------------|------------|---------------------------|--------|--------|----------|---------|
|                 |                             |            | Silt/Sand                 | Gravel | Cobble | Boulders | Rip Rap |
| 14              | Thornton Creek -mainstem    | 5          | 66                        | 22     | 1      |          | 12      |
| 14              | Matthews Creek              | 2          | 87                        | 6      |        |          | 7       |
| 14              | Unnamed trib C              | 1          | 100                       |        |        |          |         |
| 14              | Maple Creek                 | 3          | 50                        | 50     |        |          |         |
| 14              | Unnamed trib A              | 2          | 93                        | 7      |        |          |         |
| 14              | Unnamed trib B              | 1          | 40                        | 60     |        |          |         |
| 14              | Mock Creek                  | 2          | 73                        | 27     |        |          |         |
| 14              | North Branch Thornton Creek | 5          | 15                        | 63     | 17     |          | 6       |
| 14              | Littlebrook Creek           | 5          | 9                         | 45     | 46     |          |         |
| 14              | Littles Creek               | 1          | 25                        | 50     | 25     |          |         |
| 14              | South Branch Thornton Creek | 6          | 25                        | 24     | 49     | 1        | 1       |
| 14              | Meadowbrook Creek           | 2          | 100                       |        |        |          |         |
| 14              | Kramer Creek                | 1          | 92                        | 5      | 1      | 2        |         |
| 14              | Unnamed trib C              | 1          | 80                        | 20     |        |          |         |
| 14              | Willow Creek                | 5          | 39                        | 27     | 16     | 10       | 8       |
| 14              | East Fork Willow Creek      | 2          | 20                        | 40     | 40     |          |         |
| 14              | Victory Creek               | 1          | 10                        | 10     |        |          | 80      |
| 15              | Inverness Creek             | 1          | 95                        | 5      |        |          |         |
| 17              | Ravenna Creek               | 2          | 85                        | 15     |        |          |         |



TABLE 1.5.-- Stream habitat data for streams surveyed in the northeast region, August-October 2005. Locations of sample sites are displayed in Appendices 1.8-1.18. Max = maximum. Percent riffles and pools/glides are the percent of the total stream area.

| Stream system # | Sample # | Stream Name               | Date Surveyed | Distance from mouth (m) | Mean wetted width (m) | Mean depth (m) | Max depth (m) | % riffles | % pools/glides |
|-----------------|----------|---------------------------|---------------|-------------------------|-----------------------|----------------|---------------|-----------|----------------|
| 14              | 50       | Thornton Creek - mainstem | 22-Aug        | 30-80                   | 3.00                  | 0.60           | 1.20          | 0.0       | 100.0          |
| 14              | 70       | Unnamed trib C            | 29-Aug        | 0-30                    | 0.40                  | 0.02           | 0.04          | 100.0     | 0.0            |
| 14              | 72       | Maple Creek               | 29-Aug        | 450-500                 | 1.10                  | 0.08           | 0.13          | 100.0     | 0.0            |
| 14              | 71       | Unnamed trib A            | 29-Aug        | 340-390                 | 0.80                  | 0.05           | 0.06          | 100.0     | 0.0            |
| 14              | 73       | Unnamed trib A            | 29-Aug        | 740-790                 | 0.63                  | 0.05           | 0.10          | 100.0     | 0.0            |
| 14              | 74       | Unnamed trib B            | 29-Aug        | 360-410                 | 0.70                  | 0.07           | 0.20          | 100.0     | 0.0            |
| 14              | 52       | Thornton Creek - mainstem | 31-Aug        | 350-450                 | 5.17                  | 0.16           | 0.68          | 15.9      | 84.1           |
| 14              | 53       | Thornton Creek - mainstem | 30-Aug        | 500-600                 | 4.24                  | 0.20           | 0.60          | 26.9      | 73.1           |
| 14              | 54       | Thornton Creek - mainstem | 12-Sep        | 1,100-1,150             | 3.30                  | 0.33           | 0.80          | 80.8      | 19.2           |
| 14              | 76       | Mock Creek                | 15-Sep        | 460-480                 | 0.70                  | 0.03           | 0.13          | 100.0     | 0.0            |
| 14              | 77       | Mock Creek                | 15-Sep        | 770-800                 | 0.50                  | 0.03           | 0.07          | 100.0     | 0.0            |
| 14              | 56       | North Branch Thornton     | 12-Sep        | 2,850-2,901             | 2.50                  | 0.16           | 0.65          | 58.2      | 41.8           |
| 14              | 78       | Littlebrook Creek         | 8-Sep         | 0-50                    | 1.00                  | 0.20           | 0.45          | 0.0       | 100.0          |
| 14              | 79       | Littlebrook Creek         | 8-Sep         | 350-400                 | 2.00                  | 0.20           | 0.73          | 0.0       | 100.0          |
| 14              | 81       | Littlebrook Creek         | 23-Sep        | 2,350-2,375             | 0.50                  | 0.02           | 0.05          | 100.0     | 0.0            |
| 14              | 82       | Littlebrook Creek         | 23-Sep        | 2,930-2,970             | 0.50                  | 0.02           | 0.06          | 100.0     | 0.0            |
| 14              | 83       | Littles Creek             | 5-Oct         | 1,100-1,147             | 2.00                  | 0.02           | 0.26          | 93.6      | 6.4            |
| 14              | 57       | North Branch Thornton     | 17-Sep        | 3,650-3,700             | 2.62                  | 0.13           | 0.45          | 68.7      | 31.3           |
| 14              | 58       | North Branch Thornton     | 19-Sep        | 4,300-4,363             | 3.25                  | 0.19           | 0.65          | 66.2      | 33.8           |
| 14              | 59       | North Branch Thornton     | 7-Sep         | 5,150-5,200             | 2.32                  | 0.13           | 0.40          | 72.4      | 27.6           |
| 14              | 60       | North Branch Thornton     | 7-Sep         | 5,950-5,970             | 2.50                  | 0.10           | 0.20          | 100.0     | 0.0            |
| 14              | 61       | North Branch Thornton     | 7-Sep         | 6,600-6,650             | 2.28                  | 0.14           | 0.36          | 0.0       | 100.0          |
| 14              | 62       | South Branch Thornton     | 27-Sep        | 0-45                    | 0.28                  | 0.16           | 0.42          | 100.0     | 0.0            |
| 14              | 86       | Meadowbrook Creek         | 20-Sep        | 0-56                    | 0.90                  | 0.10           | 0.20          | 0.0       | 100.0          |
| 14              | 88       | Meadowbrook Pond          | 20-Sep        | 350-500                 | 4.00                  | 0.07           | 1.50          | 0.0       | 100.0          |
| 14              | 89       | Unnamed trib C            | 27-Sep        | 0-6.8                   | 0.14                  | 0.02           | 0.04          | 100.0     | 0.0            |
| 14              | 90       | Willow Creek              | 20-Sep        | 0-50.5                  | 1.48                  | 0.12           | 0.40          | 34.4      | 65.6           |
| 14              | 91       | Willow Creek              | 27-Sep        | 250-265                 | 1.00                  | 0.05           | 0.60          | 100.0     | 0.0            |
| 14              | 95       | Unnamed trib E            | 21-Sep        | 0-32                    | 1.05                  | 0.14           | 0.46          | 58.3      | 41.7           |
| 14              | 96       | Unnamed trib E            | 21-Sep        | 350-385.5               | 0.59                  | 0.06           | 0.20          | 75.4      | 24.6           |
| 14              | 92       | Willow Creek              | 21-Sep        | 350-388                 | 1.17                  | 0.09           | 0.29          | 52.8      | 47.2           |
| 14              | 93       | Willow Creek              | 21-Sep        | 920-972                 | 1.05                  | 0.13           | 0.28          | 22.5      | 77.5           |
| 14              | 94       | Willow Creek              | 21-Sep        | 1,500-1,518             | 0.90                  | 0.08           | 0.14          | 0.0       | 100.0          |
| 14              | 64       | South Branch Thornton     | 21-Sep        | 1,350-1,410             | 2.92                  | 0.27           | 0.90          | 55.5      | 44.5           |
| 14              | 65       | South Branch Thornton     | 22-Sep        | 2,250-2,297             | 2.73                  | 0.13           | 0.48          | 18.5      | 81.5           |
| 14              | 97       | Victory Creek             | 27-Sep        | 250-283                 | 0.40                  | 0.02           | 0.20          | 81.3      | 18.8           |
| 14              | 66       | South Branch Thornton     | 22-Sep        | 3,200-3,247             | 3.22                  | 0.20           | 0.42          | 21.0      | 79.0           |
| 14              | 67       | South Branch Thornton     | 22-Sep        | 3,750-3,807             | 1.93                  | 0.27           | 0.63          | 3.4       | 96.6           |
| 14              | 199      | NSCC Slough               | 1-Sep         | 4,650-4,670             | 2.20                  | 0.35           | 0.60          | 0.0       | 100.0          |
| 15              | 100      | Iverness Creek            | 28-Sep        | 0-35                    | 0.40                  | 0.03           | 0.08          | 100.0     | 0.0            |
| 17              | 105      | Ravenna Park Creek        | 4-Oct         | 175-225                 | 1.50                  | 0.10           | 0.20          | 100.0     | 0.0            |
| 17              | 106      | Ravenna Park Creek        | 5-Oct         | 600-650                 | 1.50                  | 0.08           | 0.16          | 100.0     | 0.0            |



FIGURE 1.5.-- Photo of a small pool on Matthews Creek (looking downstream), 120 m upstream from the confluence with Thornton Creek, February 2006.



FIGURE 1.6.-- Photo taken of Maple Creek looking upstream from where it flows under Sand Point Way NE.

Habitat conditions in North Branch of Thornton Creek appear to be good in most areas. Some areas are armored with rip rap. Several large, deep pools were usually present in each study section. The North Branch of Thornton Creek has two main tributaries, Littlebrook Creek and Littles Creek. The lower 1,000 m of Littlebrook Creek is daylighted, of which the lower 400 m appears to have available fish habitat. Upstream the stream is shallow with few pools (Table 1.5). The lower 500 m of Littles Creek is in culvert. The upstream daylighted reach appears to consist primarily of riffles and is relatively shallow (Table 1.5). Within Jackson Park Golf Course there are four large ponds, three adjacent to the North Branch of Thornton Creek and one on Little Creek. The average depth each pond is about 1.6 m and maximum depth is at least 2 m deep for each pond.

The South Fork of Thornton Creek is mostly daylighted from its mouth to 5<sup>th</sup> Avenue (Rm 3,807). Upstream of this the stream appears to be in a culvert until a small pond next to a Park and Ride just east of I-5, near 1<sup>st</sup> Ave and 100<sup>th</sup> St (Figure 1.7). Also there is a large pond and a small low-gradient stream on the west side of I-5 near North Seattle Community College (NSCC) that may be the headwaters of the South Branch of Thornton Creek (Appendix 1.16). The exact connection between the NSCC pond and the Park and Ride pond is unclear. Similar to the North Branch, habitat conditions in South Branch of Thornton Creek (downstream of 5<sup>th</sup> Avenue) appear to be good in most areas (Figure 1.8). Several large, deep pools were usually present in each study section. Also, LWD was present in some areas.

The other streams in the northeast region consist of Inverness Creek, Yesler Creek, and Ravenna Creek. We were unable to survey Yesler Creek, which was inaccessible. Only the upper reaches of Ravenna Creek are daylighted. The stream drains to Lake Washington through University Slough in Union Bay. We sampled two sites in Ravenna Creek; both had only riffle habitat (Table 1.5). The maximum depth was only 0.2 m and the substrate at both sites consisted of 85% sand/silt and 15% gravel (Table 1.4). Inverness Creek was substantially smaller than Ravenna Creek, it had a wetted width of 0.4 m and a maximum depth of only 0.08 m (Table 1.5).



FIGURE 1.7.-- Photo of small pond next to the Park and Ride just east of I-5, near 1<sup>st</sup> Ave and 100<sup>th</sup> St. The pond appears to be part of the upper reach of the South Branch of Thornton Creek.



FIGURE 1.8.-- Photo of Steve Damm, USFWS taking habitat measurements on the South Branch of Thornton Creek, September 2005.

*West-central region.*— All streams in the west-central region are relatively small and shallow with low streamflows and little pool habitat (Tables 1.6 and 1.7; Figures 1.9 and 1.10). Substrate was predominantly sand in each stream (Table 1.8). None of the creeks appear to be passable to fish moving upstream from Puget Sound or the Lake Washington Ship Canal. The mouth of Scheuerman Creek has a perched culvert (Figure 1.9) and Wolfe Creek goes through a long culvert before draining into Salmon Bay (Figure 1.10). In addition to stream habitat, there are a few ponds in the region (especially in the Scheuerman Creek system) that are capable of supporting fish populations.

TABLE 1.6.-- Stream habitat data for streams surveyed in the west-central region, August-September 2005. Locations of sample sites are displayed in Appendices 1.19 and 1.20. Max = maximum. Percent riffles and pools/glides are the percent of the total stream area.

| Stream system # | Sample # | Stream Name              | Date Surveyed | Distance from mouth (m) | Mean wetted width (m) | Mean depth (m) | Max depth (m) | % riffles | % pools/glides |
|-----------------|----------|--------------------------|---------------|-------------------------|-----------------------|----------------|---------------|-----------|----------------|
| 18              | 110      | Mahteen Creek            | 19-Aug        | 0-40                    | 0.40                  | 0.05           | 0.10          | 100       | 0              |
| 19              | 111      | Lawton Creek             | 17-Aug        | 50-100                  | 0.30                  | 0.02           | 0.05          | 100       | 0              |
| 20              | 112      | Wolfe Creek              | 19-Aug        | 0-50                    | 0.76                  | 0.10           | 0.20          | 100       | 0              |
| 21              | 114      | Scheuerman Creek         | 25-Aug        | 0-50                    | 1.14                  | 0.08           | 0.17          | 85.7      | 14.3           |
| 21              | 116      | Scheuerman Creek         | 19-Aug        | 280-325                 | 0.39                  | 0.05           | 0.10          | 100       | 0              |
| 21              | 117      | Scheuerman Creek         | 19-Aug        | 460-490                 | 0.60                  | 0.02           | 0.50          | 0         | 100            |
| 21              | 118      | Unnamed trib at rkm 0.28 | 19-Aug        | 0-50                    | 0.45                  | 0.05           | 0.06          | 100       | 0              |
| 22              | 122      | Owl's Creek              | 29-Sep        | 0-12                    | 0.80                  | 0.04           | 0.14          | 100       | 0              |
| 26              | 126      | Magnolia Blvd Creek      | 25-Aug        | 200-270                 | 0.55                  | 0.05           | 0.09          | 80        | 20             |

TABLE 1.7.-- Streamflow, water temperature, and conductivity measurements at various sites in Seattle streams, July-October 2005. Method is the methodology used to measure streamflow; either using a flowmeter method (F) or a volumetric measurement method (V). Measurements were usually taken sometime between 900 and 1500 h.

| Stream system # | Stream name               | Distance from mouth (m) | Method | Streamflow (cfs) | Temperature (C°) | Conductivity (µS) |
|-----------------|---------------------------|-------------------------|--------|------------------|------------------|-------------------|
| 1               | Unnamed                   | 10                      | F      | 0.120            | 10.4             | 309               |
| 4               | Broadview Creek           | 205                     | F      | 0.440            | 11.5             | 259.9             |
| 5               | Piper's Creek             | 100                     | F      | 3.630            | 12.9             | 262               |
| 8               | Blue Ridge Creek          | 350                     | F      | 0.180            | --               | --                |
| 12              | Golden Gardens Stream B   | 0                       | F      | 0.212            | 12.4             | 297.2             |
| 12              | Golden Gardens Stream E   | 20                      | F      | 0.212            | 12.5             | 230.2             |
| 14              | South Fork Thornton Creek | 0                       | F      | 1.640            | 12.7             | 250.8             |
| 14              | Unnamed trib C            | 0                       | F      | 0.029            | 12.9             | 253               |
| 14              | Willow Creek              | 0                       | F      | 0.440            | 14               | 255.1             |
| 14              | Unnamed trib E            | 0                       | F      | 0.028            | 12               | 232.9             |
| 14              | Victory Creek             | 0                       | F      | 0.113            | 15.2             | 168.4             |
| 15              | Inverness Creek           | 0                       | F      | 0.041            | 13.3             | 290.1             |
| 17              | Ravenna Creek             | 175                     | F      | 0.729            | 11.2             | 270               |
| 18              | Mahteen Creek             | 0                       | F      | 0.095            | --               | --                |
| 19              | Lawton Creek              | 25                      | F      | 0.247            | 13.9             | 271.6             |
| 20              | Wolfe Creek               | 0                       | F      | 0.050            | 14.9             | 261.6             |
| 21              | Scheuerman Creek          | 0                       | V      | 0.176            | 12.2             | 310               |
| 22              | Owl's Creek               | 0                       | F      | 0.025            | 12.8             | 293.7             |
| 26              | Magnolia Blvd Creek       | 200                     | V      | 0.055            | 15               | 262.3             |
| 27              | Washington Park Creek     | 105                     | V      | 0.026            | 15.6             | 156.7             |
| 29              | Interlaken Creek -reach C | 0                       | V      | 0.0045           | 15.5             | 178.5             |
| 30              | Boren Park Creek          | 0                       | F      | 0.018            | 13.4             | 214.5             |
| 31              | Madrona Park North Trib   | 275                     | V      | 0.0031           | --               | --                |
| 32              | Madrona Park South Trib   | 50                      | V      | 0.015            | --               | --                |
| 35              | Puget Creek               | 0                       | V      | 0.0057           | --               | --                |
| 35              | Unnamed trib A            | 7                       | V      | 0.0035           | --               | --                |
| 35              | Unnamed trib B            | 14                      | V      | 0.0069           | --               | --                |
| 38              | Schmitz Creek             | 0                       | F      | 0.269            | 14               | 264.8             |
| 38              | Unnamed trib G            | 15                      | V      | 0.034            | 14               | 228.5             |
| 39              | Mee-Kwa-Mooks Creek       | 130                     | V      | 0.061            | 13.4             | 336               |
| 40              | Pelly Creek               | 130                     | V      | 0.0016           | 14.7             | 224.1             |
| 42              | Fauntleroy Creek          |                         | F      | 0.650            | 14.3             | 255               |
| 44              | Durham Creek              | 150                     | F      | 0.500            | 13.4             | 305               |
| 49              | Taylor Creek              | 0                       | F      | 0.382            | --               | --                |
| 49              | East Fork Taylor Creek    | 0                       | F      | 0.012            | --               | --                |
| 49              | West Fork Taylor Creek    | 0                       | F      | 0.480            | --               | --                |

TABLE 1.8.-- Total combined substrate scores for streams in the west-central region.

| Stream system # | Stream name      | # of Sites | Overall Substrate Score % |        |        |                  |
|-----------------|------------------|------------|---------------------------|--------|--------|------------------|
|                 |                  |            | Silt/Sand                 | Gravel | Cobble | Boulders Rip Rap |
| 18              | Mahteen Creek    | 1          | 100                       |        |        |                  |
| 19              | Lawton Creek     | 1          | 100                       |        |        |                  |
| 20              | Wolf Creek       | 2          | 100                       |        |        |                  |
| 21              | Scheuerman Creek | 4          | 71                        | 19     | 7      | 3                |
| 21              | Unnamed trib A   | 1          | 99                        | 1      |        |                  |
| 22              | Owl's Creek      | 2          | 85                        | 15     |        |                  |
| 26              | West Magnolia    | 1          | 95                        | 5      |        |                  |



FIGURE 1.9.-- Photos of Wolfe Creek (stream system #20). The left photo was taken at the downstream end of the daylighted section where it enters into a culvert (sample location #112). The right photo shows the overall shallow depth of this stream at this site and the predominantly sand substrate.



FIGURE 1.10.-- Photos of Scheuerman Creek (stream system #21). The left photo was taken at the creek mouth where it exits a culvert and then drops sharply to Puget Sound. The right photo of Scheuerman Creek was taken approximately 30 m upstream of the culvert and shows the overall shallow depth of this stream.

East-central region.— The largest stream in this region is Washington Park Creek. The stream has a perched culvert at Rm 105 which most likely restricts upstream movements of fish. The stream is approximately 1,650 m long and has some pool habitat (Table 1.9) and some riparian vegetation that could provide cover for fish; however the stream is relatively small with little streamflow (0.026 cfs) and predominantly sand/silt substrate (Table 1.10). At the upper end of this system there are two ponds, the koi pond in the Japanese Gardens and a small pond on a tributary at Rm 1,500. The streams in the Interlaken, Boren, and Madrona Park are quite small and shallow with steep gradients. Frink Creek had more streamflow but was still a small, shallow stream.



TABLE 1.9.-- Stream habitat data for streams surveyed in the east-central region, July-August 2005. Locations of sample sites are displayed in Appendices 1.21 and 1.22. Max = maximum; dashes indicate that no data was collected. Percent riffles and pools/glides are the percent of the total stream area.

| Stream system # | Sample # | Stream Name               | Date Surveyed | Distance from mouth (m) | Mean wetted width (m) | Mean depth (m) | Max depth (m) | % riffles | % pools/glides |
|-----------------|----------|---------------------------|---------------|-------------------------|-----------------------|----------------|---------------|-----------|----------------|
| 18              | 130      | Washington Park Creek     | 11-Aug        | 0-50                    | 0.85                  | 0.12           | 0.50          | 34.0      | 66.0           |
| 19              | 230      | Washington Park Creek     | 11-Aug        | 50-105                  | 0.85                  | 0.14           | 0.47          | 64.7      | 35.3           |
| 20              | 131      | Washington Park Creek     | 11-Aug        | 850-900                 | 0.77                  | 0.06           | 0.19          | 82.0      | 18.0           |
| 28              | 134      | Interlaken Area - reach A | 17-Aug        | 80-90                   | 0.10                  | --             | --            | --        | --             |
| 29              | 135      | Interlaken Area - reach B | 19-Aug        | 0-20                    | 0.15                  | 0.01           | 0.01          | 100       | 0              |
| 29              | 136      | Interlaken Area - reach C | 19-Aug        | 0-50                    | 0.10                  | 0.01           | 0.50          | 100       | 0              |
| 30              | 137      | Boren Park Area           | 19-Aug        | 0-30                    | 0.50                  | 0.03           | 0.08          | 100       | 0              |
| 31              | 138      | Madrona Park N. Trib      | 10-Aug        | 275                     | 0.40                  | 0.01           | 0.02          | 100       | 0              |
| 31              | 139      | Madrona Park S. Trib      | 10-Aug        | 0                       | 0.45                  | 0.05           | 0.15          | 100       | 0              |
| 33              | 140      | Frink Cr                  | 28-Jul        | 575-625                 | 0.20                  | 0.03           | 0.20          | 100       | 0              |

TABLE 1.10.-- Total combined substrate scores for streams in the east-central region.

| Stream system # | Stream name                      | # of Sites | Overall Substrate Score % |        |        |          |         |
|-----------------|----------------------------------|------------|---------------------------|--------|--------|----------|---------|
|                 |                                  |            | Silt/Sand                 | Gravel | Cobble | Boulders | Rip Rap |
| 27              | Washington Park Creek            | 4          | 85                        | 6      | 9      |          |         |
| 28              | Interlaken Creek (reach A and B) | 2          | 100                       |        |        |          |         |
| 29              | Interlaken Creek - reach C       | 1          | 90                        |        |        |          | 10      |
| 30              | Boren Park Creek                 | 1          | 100                       |        |        |          |         |
| 31              | Madrona Park North Trib          | 2          | 100                       |        |        |          |         |
| 32              | Madrona Park South Trib          | 1          | 25                        |        |        |          | 75      |
| 33              | Frink Creek                      | 2          | 75                        | 20     | 5      |          |         |

*Southwest region.*— The largest stream in the southwest region is Longfellow Creek. We surveyed a total of 10 sites on the mainstem of Longfellow Creek. Habitat conditions appear to be generally good throughout the 5.7 km that we surveyed. Large, deep pools are common. Of the daylighted sections of Puget Creek, only the lower 60 m had any water during the summer. Upstream the streambed was dry. Most of the streamflow in Puget Creek is from a right-bank tributary at Rm 54 (Figure 1.11; Table 1.7). The best fish habitat in the Puget Creek system is the lower 14 m of this tributary which includes a plunge pool at the base of a culvert (Figure 1.11). We did not assess habitat above the culvert but the gradient is steep and it's doubtful if

there is any available habitat. Another small tributary to Puget Creek at Rm 12 also adds some streamflow.



FIGURE 1.11.-- Photo taken of the lower 13 m of a right-bank tributary to Puget Creek. The photo was taken on February 8, 2006. Most of the streamflow in Puget Creek comes from this tributary.

The lower section of Schmitz Creek is in a culvert, which appears to be a complete barrier to upstream fish movements. Upstream of the culvert, the stream is located in forested park. The lower 700 m of the daylighted reach is about 1.5-m wide with only a few small, shallow pools (Table 1.11) and little woody debris. The substrate is predominantly sand (Table 1.12). Overall, Schmitz Creek appears to have limited available fish habitat. Fauntleroy Creek is the second largest stream in the southwest region. Upstream of the lower culvert, several pieces of large woody debris have been added as part of a restoration effort. The amount of pool habitat decreases in upstream areas. The substrate is predominantly sand throughout the stream.

The furthest downstream section of Durham Creek (stream system #44) that is daylighted is located in the South Park Pea Patch. This stream section has a low gradient and has adequate pool habitat and cover due to large amounts of water cress *Nasturtium officinale*. Just upstream of this section is a steep culvert which may restrict upstream movements of fish. Upstream the stream is in a culvert under Highway 509. Between Myers Way S and Highway 509 there are

three streams, it appears the middle stream is directly connected to Durham Creek but it is unclear if the other streams (stream systems #'s 45 and 46) are directly connected to Durham Creek. Although these streams have adequate habitat and streamflow, they appear to be above impassable culverts.

TABLE 1.11.-- Stream habitat data for streams surveyed in the southwest region, July-October 2005. Locations of sample sites are displayed in Appendices 1.23-1.28. Max = maximum; dashes indicate that no data was collected. Percent riffles and pools/glides are the percent of the total stream area.

| Stream system # | Sample # | Stream Name         | Date Surveyed | Distance from mouth (m) | Mean wetted width (m) | Mean depth (m) | Max depth (m) | % riffles | % pools/glides |
|-----------------|----------|---------------------|---------------|-------------------------|-----------------------|----------------|---------------|-----------|----------------|
| 34              | 150      | Longfellow Creek    | 1-Aug         | 0-80                    | 2.36                  | 0.17           | 0.70          | 31.0      | 69.0           |
| 34              | 146      | Golf Course trib    | 1-Aug         | 0-27                    | 0.30                  | 0.10           | 0.20          | 0.0       | 100.0          |
| 34              | 147      | Unnamed trib A      | 1-Aug         | 225-245                 | 0.10                  | 0.05           | 0.10          | --        | --             |
| 34              | 151      | Longfellow Creek    | 1-Aug         | 1,000-1,050             | 3.00                  | 0.50           | 1.00          | 0.0       | 100.0          |
| 34              | 152      | Longfellow Creek    | 1-Aug         | 1,650-1,685             | 1.78                  | 0.13           | 0.35          | 76.2      | 23.8           |
| 34              | 153      | Longfellow Creek    | 2-Aug         | 2,150-2,220             | 2.13                  | 0.15           | 0.30          | 63.8      | 36.2           |
| 34              | 154      | Longfellow Creek    | 2-Aug         | 2,850-2,917             | 2.23                  | 0.15           | 0.55          | 42.4      | 57.6           |
| 34              | 155      | Longfellow Creek    | 2-Aug         | 3,400-3,450             | 4.00                  | 0.25           | 0.65          | 0.0       | 100.0          |
| 34              | 156      | Longfellow Creek    | 2-Aug         | 3,775-3,822             | 2.10                  | 0.09           | 0.90          | 89.0      | 11.0           |
| 34              | 148      | Unnamed trib B      | 2-Aug         | 0-10                    | 0.10                  | 0.02           | 0.05          | --        | --             |
| 34              | 149      | Unnamed trib C      | 3-Aug         | 0-3                     | 0.50                  | 0.01           | 0.05          | 0.0       | 100.0          |
| 34              | 157      | Longfellow Creek    | 3-Aug         | 4,850-4,898             | 1.67                  | 0.12           | 0.85          | 67.5      | 32.5           |
| 34              | 158      | Longfellow Creek    | 3-Aug         | 5,300-5,347             | 1.78                  | 0.15           | 0.50          | 39.5      | 60.5           |
| 34              | 159      | Longfellow Creek    | 4-Aug         | 5,600-5,647             | 1.51                  | 0.14           | 0.45          | 51.1      | 48.9           |
| 35              | 160      | Puget Creek         | 9-Aug         | 0-54                    | 0.73                  | 0.07           | 0.27          | 51.7      | 48.3           |
| 35              | 161      | Unnamed trib A      | 9-Aug         | 0-6.5                   | 0.30                  | 0.05           | 0.10          | 100.0     | 0.0            |
| 35              | 162      | Unnamed trib B      | 9-Aug         | 0-13.5                  | 1.29                  | 0.16           | 0.61          | 35.5      | 64.5           |
| 37              | 164      | Fairmount Creek     | 28-Jul        | 275-280                 | 0.10                  | 0.04           | 0.20          | 100.0     | 0.0            |
| 38              | 165      | Schmitz Creek       | 10-Aug        | 0-50                    | 1.50                  | 0.10           | 0.14          | 100.0     | 0.0            |
| 38              | 166      | Schmitz Creek       | 10-Aug        | 250-300                 | 1.50                  | 0.10           | 0.27          | 80.0      | 20.0           |
| 38              | 167      | Schmitz Creek       | 10-Aug        | 575-625                 | 1.50                  | 0.10           | 0.20          | 100.0     | 0.0            |
| 38              | 170      | Unnamed trib G      | 10-Aug        | 0-20                    | 0.40                  | 0.03           | 0.06          | 100.0     | 0.0            |
| 38              | 168      | Schmitz Creek       | 10-Sep        | 675-710                 | 1.50                  | 0.03           | 0.07          | 100.0     | 0.0            |
| 38              | 171      | Unnamed trib K      | 10-Aug        | 85-100                  | 0.40                  | 0.04           | 0.07          | 100.0     | 0.0            |
| 38              | 169      | Schmitz Creek       | 10-Aug        | 710-725                 | 0.70                  | 0.04           | 0.07          | 100.0     | 0.0            |
| 39              | 172      | Mee-Kwa-Mooks Creek | 10-Aug        | 130-140                 | 0.28                  | 0.03           | 0.10          | 100.0     | 0.0            |
| 39              | 272      | Unnamed trib A      | 10-Aug        | 0-10                    | 0.20                  | 0.015          | 0.04          | 100.0     | 0.0            |
| 40              | 173      | Pelly Creek         | 10-Aug        | 130-140                 | 0.30                  | 0.02           | 0.08          | 100.0     | 0.0            |
| 42              | 180      | Fauntleroy Creek    | 18-Aug        | 0-85                    | 2.88                  | 0.05           | 0.50          | 42.9      | 57.1           |
| 42              | 181      | Fauntleroy Creek    | 18-Oct        | 250-353                 | --                    | --             | 0.46          | --        | --             |
| 42              | 183      | Fauntleroy Creek    | 18-Aug        | 725-775                 | 1.50                  | 0.06           | 0.25          | 85.0      | 15.0           |
| 42              | 184      | Unnamed trib A      | 18-Aug        | 0-20                    | --                    | --             | --            | 100.0     | 0.0            |
| 44              | 190      | Durham Creek        | 9-Aug         | 0-78                    | 2.53                  | --             | 0.53          | 0.0       | 100.0          |
| 44              | 191      | Durham Creek        | 9-Aug         | 150-200                 | 1.86                  | 0.19           | 0.42          | 90.9      | 9.1            |
| 45              | 193      | Durham Creek        | 2-Aug         | 450-487                 | 0.90                  | 0.05           | 0.34          | 100.0     | 0.0            |
| 45              | 194      | Durham Creek B      | 2-Aug         | 0-10                    | 1.00                  | 0.06           | 0.20          | 100.0     | 0.0            |
| 46              | 192      | Durham Creek C      | 2-Aug         | 0-50                    | 1.50                  | 0.20           | 0.50          | 0.0       | 100.0          |

TABLE 1.12.-- Total combined substrate scores for streams in the southwest region.

| Stream system # | Stream Name          | # of sites | Overall Substrate Score % |        |        |          |         |
|-----------------|----------------------|------------|---------------------------|--------|--------|----------|---------|
|                 |                      |            | Silt/Sand                 | Gravel | Cobble | Boulders | Rip Rap |
| 35              | Puget Creek          | 1          | 10                        | 70     | 20     |          |         |
| 35              | Unnamed trib A       | 1          |                           | 19     | 81     |          |         |
| 37              | Fairmount Creek      | 1          | 100                       |        |        |          |         |
| 38              | Schmitz Creek        | 4          | 82                        | 18     |        |          |         |
| 38              | Schmitz Creek Trib G | 1          | 100                       |        |        |          |         |
| 39              | Mee-Kwa-Mooks Creek  | 1          | 95                        | 5      |        |          |         |
| 40              | Pelly Creek          | 1          | 100                       |        |        |          |         |
| 42              | Fauntleroy Creek     | 4          | 84                        | 14     | 2      |          |         |
| 42              | Unnamed trib A       | 1          | 100                       |        |        |          |         |
| 44              | Durham Creek         | 3          | 87                        | 7      | 6      |          |         |
| 45              | Durham Creek B       | 1          | 45                        | 8      | 35     | 12       |         |
| 46              | Durham Creek C       | 1          | 100                       |        |        |          |         |

*Southeast region.*— The southeast region consists of only three stream systems: Mt. Baker Creek, Mapes Creek, and Taylor Creek. The lower sections of both Mt. Baker Creek and Mapes Creek are in culverts. The daylighted section of Mt. Baker Creek is only about 100-m long and is relatively small; however, there are three man-made barriers that create three small pools (each roughly 13-m long by 3.5-m wide). We surveyed 38 m upstream of the pools and the stream is only 0.1 m wide (Table 1.13). Mapes Creek is a substantially larger stream system, with the lower 900 m of the stream in a culvert. The daylighted section of Mapes Creek is generally small with mostly sand substrate (Table 1.14). The stream also includes three ponds (total length, 82 m) in the Kabota Gardens.

Taylor Creek is the largest stream in the southeast region. An impassable culvert under Rainier Avenue limits upstream movement of fish in Taylor Creek. Habitat conditions in the lower 500 m are generally good and includes good pool habitat. Upstream, pools are infrequent and those that are present are usually small and shallow. In the headwaters, the stream splits into two forks (East Fork and West Fork). Most of the summer streamflow is from the West Fork (Table 1.7).

TABLE 1.13.-- Stream habitat data for streams surveyed in the southeast, June-July 2005. Locations of sample sites are displayed in Appendices 1.29 and 1.30. Max = maximum. Percent riffles and pools/glides are the percent of the total stream area.

| Stream system # | Sample # | Stream Name            | Date Surveyed | Distance from mouth (m) | Mean wetted width (m) | Mean depth (m) | Max depth (m) | % riffles | % pools/glides |
|-----------------|----------|------------------------|---------------|-------------------------|-----------------------|----------------|---------------|-----------|----------------|
| 47              | 200      | Mt. Baker              | 25-Jul        | 0-50                    | 1.82                  | 0.31           | 0.55          | 7.1       | 92.9           |
| 48              | 201      | Mapes Creek            | 25-Jul        | 900-935                 | 1.10                  | 0.16           | 0.40          | 41.6      | 58.4           |
| 49              | 203      | Taylor Creek           | 30-Jun        | 0-50                    | 2.48                  | 0.09           | 0.30          | 29.0      | 71.0           |
| 49              | 205      | Taylor Creek           | 30-Jun        | 825-875                 | 1.75                  | 0.10           | 0.25          | 100.0     | 0.0            |
| 49              | 206      | East Fork Taylor Creek | 7-Jul         | 0-50                    | 1.24                  | 0.03           | 0.25          | 96.8      | 3.2            |
| 49              | 207      | East Fork Taylor Creek | 19-Jul        | 50-100                  | 1.27                  | 0.09           | 0.25          | 83.2      | 16.8           |

TABLE 1.14.-- Total combined substrate scores for streams in the southeast region.

| Stream system # | Stream Name            | # of sites | Overall Substrate Score % |        |        |          |         |
|-----------------|------------------------|------------|---------------------------|--------|--------|----------|---------|
|                 |                        |            | Silt/Sand                 | Gravel | Cobble | Boulders | Rip Rap |
| 47              | Mt. Baker Park Creek   | 1          | 84                        | 6      | 10     |          |         |
| 48              | Mapes Creek            | 2          | 98                        | 1      | 1      |          |         |
| 49              | Taylor Creek           | 3          | 39                        | 39     | 18     | 4        |         |
| 49              | East Fork Taylor Creek | 2          | 57                        | 35     | 8      |          |         |
| 49              | West Fork Taylor Creek | 1          | 50                        | 30     | 15     | 5        |         |

## CHAPTER 2. FISH DISTRIBUTION SURVEYS AND STREAM TYPING

### Introduction and Methods

*Fish distribution surveys*.-- We surveyed each stream in Seattle to determine the distribution of each fish species (salmonids and non-salmonids). Results of these fish surveys as well as habitat surveys (Chapter 1) were then use to classify each stream. Fish surveys were primarily conducted during the summer low-flow period of June to October 2005. We also sampled select locations during winter-flow conditions in February of 2006 to determine if the fish distribution had expanded to more upstream areas. In areas that we detected a range expansion, we surveyed again during summer low-flow period in August 2006 to determine if conditions between summer 2005 and 2006 were similar. Sampling sites were determined with the aid of maps and aerial photos of each watershed. We used a systematic sampling scheme to survey each watershed. Sample stream sections were generally 300 to 500 m apart for each stream, except in Thornton Creek where sample sections were 800 to 1,000 m apart. The first sample section began at the mouth of the stream, or as in many cases, where the stream enters a culvert that then transports the stream to Puget Sound, the Duwamish River, or Lake Washington. The exact location of each survey sample section was often based on accessibility and recognizable landmarks (i.e., bridges or culverts). We attempted to locate areas of a stream where sample section lengths could be at least 50 m in length and incorporate at least two habitat types (pools, riffles, or glides). This was not always the case and sample sections were sometimes less than 50 m in length with only a single habitat type. Supplemental sample sections were also surveyed at some areas to sample high quality habitat (i.e., deep pools) or more precisely document the distribution of each cottid species in the lower reaches of major creeks.

Our primary method of sampling consisted of backpack electrofishing using a Smith-Root LR-24® electrofisher system. Backpack electrofishing was generally conducted in an upstream direction with one or more individuals following behind or along side the electrofisher operator to collect the stunned fish. Fish that were stunned by the electrical field were removed from the stream with long handle dip nets and placed in a

recovery bucket. Additional surveys were conducted in September 2006 with a lamprey electrofishing unit (University of Wisconsin model #ABP-2) to attempt to find additional lamprey sites. We also used, when necessary, beach seines or gill nets to collect fish. Beach seines were used in ponds or deep waters of slow moving streams. The beach seine was 9.1-m long and 1.8-m deep, with a 1.8-m deep by 1.8-m long bag in the center. The mesh size in the wings was 6-mm stretch mesh, while the bag was 2-mm stretch mesh. We also used gill nets to sample some ponds for salmonids. Gill nets were between 6.5 and 12.5 m in length, 2.0 m deep with 2-cm square mesh. Nets were deployed perpendicular to shore and secured on both banks. Gill nets were deployed at dusk and removed just after sunrise. We also conducted snorkel surveys at a few pond sites to survey a large area. Snorkel surveys were conducted at night and were only done during the winter.

After sampling was completed at each reach, captured fish were placed in an anesthetizing water bath of MS-222. Fish were identified and the length (nearest mm) was measured. After fish had recovered, they were then placed back into the stream from the habitat that they were captured from. Often a digital photo was taken of the fish. Unidentifiable species were retained for identification in the laboratory.

*Stream typing.*-- All streams that were surveyed for fish (this chapter) and habitat (Chapter 1) were classified according to the Washington Department of Natural Resources interim and permanent stream typing system (WAC 222-16-031) (Table 2.1). The interim and permanent water typing methods relies on collecting both physical habitat parameters and fish presence/absence to obtain a water typing classification. This classification is then represented by a number (interim), or letter / letter combination (permanent), that is then given to indicate the stream type. Since this stream typing classification scheme was developed for more forested, less developed areas, it is often difficult to assign typing for streams in a heavily urbanized areas.

Table 2.1.-- Washington Department of Natural Resources water typing conversion table.

| Interim Water Typing | Permanent Water Typing |
|----------------------|------------------------|
| Type 1               | Type "S"               |
| Type 2 and 3         | Type "F"               |
| Type 4               | Type "Np"              |
| Type 5               | Type "Ns"              |

Definitions of each stream typing system and classification are as follows:

***Interim Water Typing Definitions***

**Type 1 Water**, means all waters, within their ordinary high-water mark, as inventoried as "shorelines of the state" under chapter 90.58 RCW and the rules promulgated pursuant to chapter 90.58 RCW, but not including those waters' associated wetlands as defined in chapter 90.58 RCW.

**Type 2 Water**, means segments of natural waters which are not classified as Type 1 Water and have a high fish, wildlife, or human use. These are segments of natural waters and periodically inundated areas of their associated wetlands, which:

(a) Are diverted for domestic use by more than 100 residential or camping units or by a public accommodation facility licensed to serve more than 10 persons, where such diversion is determined by the department to be a valid appropriation of water and only considered Type 2 Water upstream from the point of such diversion for 1,500 feet or until the drainage area is reduced by 50 percent, whichever is less;

(b) Are diverted for use by federal, state, tribal or private fish hatcheries. Such waters shall be considered Type 2 Water upstream from the point of diversion for 1,500 feet, including tributaries if highly significant for protection of downstream water quality. The department may allow additional harvest beyond the requirements of Type 2 Water designation provided by the department of fish and wildlife, department of ecology, the affected tribes and interested parties that:

(i) The management practices proposed by the landowner will adequately protect water quality for the fish hatchery; and

(ii) Such additional harvest meets the requirements of the water type designation that would apply in the absence of the hatchery;

(c) Are within a federal, state, local or private campground having more than 30 camping units: Provided, That the water shall not be considered to enter a campground until it reaches the boundary of the park lands available for public use and comes within 100 feet of a camping unit.

(d) Are used by fish for spawning, rearing or migration. Waters having the following characteristics are presumed to have highly significant fish populations:

(i) Stream segments having a defined channel 20 feet or greater within the bankfull width and having a gradient of less than 4 percent.

(ii) Lakes, ponds, or impoundments having a surface area of 1 acre or greater at seasonal low water; or

(e) Are used by fish for off-channel habitat. These areas are critical to the maintenance of optimum survival of fish. This habitat shall be identified based on the following criteria:

(i) The site must be connected to a fish bearing stream and be accessible during some period of the year; and

(ii) The off-channel water must be accessible to fish through a drainage with less than a 5% gradient.

**Type 3 Water**, means segments of natural waters which are not classified as Type 1 or 2 Waters and have a moderate to slight fish, wildlife, or human use. These are segments of natural waters and periodically inundated areas of their associated wetlands which:

(a) Are diverted for domestic use by more than 10 residential or camping units or by a public accommodation facility licensed to serve more than 10 persons, where such diversion is determined by the department to be a valid appropriation of water and the only practical water source for such users. Such waters shall be considered to be Type 3 Water upstream from the point of such diversion for 1,500 feet or until the drainage area is reduced by 50 percent, whichever is less;

(b) Are used by fish for spawning, rearing or migration. The requirements for determining fish use are described in the board manual section 13. If fish use has not been determined:

(i) Waters having any of the following characteristics are presumed to have fish use:

(A) Stream segments having a defined channel of 2 feet or greater within the bankfull width in Western Washington; or 3 feet or greater in width in Eastern Washington; and having a gradient of 16 percent or less;

(B) Stream segments having a defined channel of 2 feet or greater within the bankfull width in Western Washington; or 3 feet or greater within the bankfull width in Eastern Washington, and having a gradient greater than 16 percent and less than or equal to 20 percent, and having greater than 50 acres in contributing basin size in Western Washington or greater than 175 acres contributing basin size in Eastern Washington, based on hydrographic boundaries;



(C) Ponds or impoundments having a surface area of less than 1 acre at seasonal low water and having an outlet to a fish stream;

(D) Ponds or impoundments having a surface area greater than 0.5 acre at seasonal low water.

(ii) The department shall waive or modify the characteristics in (i) of this subsection where:

(A) Waters have confirmed, long term, naturally occurring water quality parameters incapable of supporting fish;

(B) Snowmelt streams have short flow cycles that do not support successful life history phases of fish. These streams typically have no flow in the winter months and discontinue flow by June 1; or

(C) Sufficient information about a geomorphic region is available to support a departure from the characteristics in (i) of this subsection, as determined in consultation with the department of fish and wildlife, department of ecology, affected tribes and interested parties.

**Type 4 Water**, means all segments of natural waters within the bankfull width of defined channels that are perennial nonfish habitat streams. Perennial streams are flowing waters that do not go dry any time of a year of normal rainfall and include the intermittent dry portions of the perennial channel below the uppermost point of perennial flow.

**Type 5 Waters**, means all segments of natural waters within the bankfull width of the defined channels that are not Type 1, 2, 3, or 4 Waters. These are seasonal, nonfish habitat streams in which surface flow is not present for at least some portion of the year and are not located downstream from any stream reach that is a Type 4 Water. Type 5 Waters must be physically connected by an above-ground channel system to Type 1, 2, 3, or 4 Waters.

For purposes of this section:

(a) "Residential unit" means a home, apartment, residential condominium unit or mobile home, serving as the principal place of residence.

(b) "Camping unit" means an area intended and used for:

(i) Overnight camping or picnicking by the public containing at least a fireplace, picnic table and access to water and sanitary facilities; or

(ii) A permanent home or condominium unit or mobile home not qualifying as a "residential unit" because of part time occupancy.

(c) "Public accommodation facility" means a business establishment open to and licensed to serve the public, such as a restaurant, tavern, motel or hotel.

(d) "Natural waters" only excludes water conveyance systems which are artificially constructed and actively maintained for irrigation.

(e) "Seasonal low flow" and "seasonal low water" mean the conditions of the 7-day, 2-year low water situation, as measured or estimated by accepted hydrologic techniques recognized by the department.

(f) "Channel width and gradient" means a measurement over a representative section of at least 500 linear feet with at least 10 evenly spaced measurement points along the normal stream channel but excluding unusually wide areas of negligible gradient such as marshy or swampy areas, beaver ponds and impoundments. Channel gradient may be determined utilizing stream profiles plotted from United States geological survey topographic maps (See board manual section 23).

### ***Permanent Water Typing Definitions***

**Type S Water** means all waters, within their bankfull width, as inventoried as "shorelines of the state" under chapter [90.58 RCW](#) and the rules promulgated pursuant to chapter [90.58 RCW](#) including periodically inundated areas of their associated wetlands.

**Type F Water** means segments of natural waters other than Type S Waters, which are within the bankfull widths of defined channels and periodically inundated areas of their associated wetlands, or within lakes, ponds, or impoundments having a surface area of 0.5 acre or greater at seasonal low water and which in any case contain fish habitat or are described by one of the following four categories:

(a) Waters, which are diverted for domestic use by more than 10 residential or camping units or by a public accommodation facility licensed to serve more than 10 persons, where such diversion is determined by the department to be a valid appropriation of water and the only practical water source for such users. Such waters shall be considered to be Type F Water upstream from the point of such diversion for 1,500 feet or until the drainage area is reduced by 50 percent, whichever is less;

(b) Waters, which are diverted for use by federal, state, tribal or private fish hatcheries. Such waters shall be considered Type F Water upstream from the point of diversion for 1,500 feet, including tributaries if

highly significant for protection of downstream water quality. The department may allow additional harvest beyond the requirements of Type F Water designation provided the department determines after a landowner-requested on-site assessment by the department of fish and wildlife, department of ecology, the affected tribes and interested parties that:

(i) The management practices proposed by the landowner will adequately protect water quality for the fish hatchery; and

(ii) Such additional harvest meets the requirements of the water type designation that would apply in the absence of the hatchery;

(c) Waters, which are within a federal, state, local, or private campground having more than 10 camping units: Provided, That the water shall not be considered to enter a campground until it reaches the boundary of the park lands available for public use and comes within 100 feet of a camping unit, trail or other park improvement;

(d) Riverine ponds, wall-based channels, and other channel features that are used by fish for off-channel habitat. These areas are critical to the maintenance of optimum survival of fish. This habitat shall be identified based on the following criteria:

(i) The site must be connected to a fish habitat stream and accessible during some period of the year; and

(ii) The off-channel water must be accessible to fish.

**Type Np Water** means all segments of natural waters within the bankfull width of defined channels that are perennial nonfish habitat streams. Perennial streams are flowing waters that do not go dry any time of a year of normal rainfall and include the intermittent dry portions of the perennial channel below the uppermost point of perennial flow.

**Type Ns Water** means all segments of natural waters within the bankfull width of the defined channels that are not Type S, F, or Np Waters. These are seasonal, nonfish habitat streams in which surface flow is not present for at least some portion of a year of normal rainfall and are not located downstream from any stream reach that is a Type Np Water. Ns Waters must be physically connected by an above-ground channel system to Type S, F, or Np Waters.

For purposes of this section:

(a) "Residential unit" means a home, apartment, residential condominium unit or mobile home, serving as the principal place of residence.

(b) "Camping unit" means an area intended and used for:

(i) Overnight camping or picnicking by the public containing at least a fireplace, picnic table and access to water and sanitary facilities; or

(ii) A permanent home or condominium unit or mobile home not qualifying as a "residential unit" because of part time occupancy.

(c) "Public accommodation facility" means a business establishment open to and licensed to serve the public, such as a restaurant, tavern, motel or hotel.

(d) "Natural waters" only excludes water conveyance systems which are artificially constructed and actively maintained for irrigation.

(e) "Seasonal low flow" and "seasonal low water" mean the conditions of the 7-day, 2-year low water situation, as measured or estimated by accepted hydrologic techniques recognized by the department.

(f) "Channel width and gradient" means a measurement over a representative section of at least 500 linear feet with at least 10 evenly spaced measurement points along the normal stream channel but excluding unusually wide areas of negligible gradient such as marshy or swampy areas, beaver ponds and impoundments. Channel gradient may be determined utilizing stream profiles plotted from United States geological survey topographic maps (see board manual section 23).

(g) "Intermittent streams" means those segments of streams that normally go dry.

(h) "Fish habitat" means habitat which is used by any fish at any life stage at any time of the year, including potential habitat likely to be used by fish which could be recovered by restoration or management and includes off-channel habitat.

## Results

During the summer low-flow period (June-October 2005), we surveyed a total of 167 sample sections throughout the 49 watersheds. A total of 27 sample sections were resurveyed during the winter.

### *Fish Distribution by Region*

Northwest region.-- The Piper's Creek system was the only system in the northwest region in which we captured fish. In our survey of Piper's Creek, we found cottids upstream to Rm 200, coho salmon up to the confluence of Venema Creek at Rm 620, and cutthroat trout up to Rm 1,850. During the summer 2005 surveys, we found the cutthroat trout distribution only extended upstream to two large culverts (twin culverts) at Rm 1,750 (Table 2.2). During the winter 2006 surveys, we collected three cutthroat trout 100 m upstream of the twin culverts and two cutthroat trout were collected from the same location in August 2006 (Table 2.3).

In Venema Creek, we found cutthroat trout from the mouth to a large log jam located at Rm 320. We sampled a few sites (including a few spot samples) upstream of the log jam during the summer of 2005 and winter of 2006 but did not capture any fish. We sampled Mohlendorph Creek during the summer of 2005, but there was little streamflow, and no fish were captured. Mohlendorph Creek was re-sampled during the winter of 2006 and we captured two juvenile cutthroat trout at the mouth below the cement weir and another juvenile cutthroat trout 50 m upstream of the mouth. These were the only three fish we captured during all of our surveys on Mohlendorph Creek.

We also captured cutthroat trout on two other tributaries to Piper's Creek. Three cutthroat trout were collected in the lower 20 m of a right-bank tributary located at Rm 970 (called North Creek in Thomas 1992). In another right-bank tributary at Rm 1,450 (called Viewlands Creek in Thomas 1992), four cutthroat trout were captured.

TABLE 2.2.-- Number of fish collected in the Northwest region watersheds during fish distribution surveys, October 2005. NS = no sample. Gear used was either electrofishing (E), beach seine (B), or not fished (NF, only habitat information collected). Locations of sample sites are displayed in Appendices 1.1-1.7. Fish caught of sample #'s greater than 500 are results from the first pass of reference site sampling (Chapter 3). Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are coho salmon (COH), cutthroat trout (CUT), coastrange sculpin (CRS), prickly sculpin (PKS), staghorn sculpin (SHS), and shorthose sculpin (SNS).

| Stream system # | Sample # | Stream name             | Date Sampled | Distance from mouth (m) | Gear | Stream Type | Number caught |     |     |     |     |     |
|-----------------|----------|-------------------------|--------------|-------------------------|------|-------------|---------------|-----|-----|-----|-----|-----|
|                 |          |                         |              |                         |      |             | COH           | CUT | CRS | PKS | SHS | SNS |
| 1               | 1        | Unnamed                 | 4-Oct        | 10                      | NF   | 4           |               |     |     |     |     |     |
| 1               | 2        | Unnamed                 | 4-Oct        | 120-147                 | B    | 4           |               |     |     |     |     |     |
| 1               | 3        | Unnamed                 | 4-Oct        | 170-216                 | E    | 4           |               |     |     |     |     |     |
| 2               | NS       | Unnamed                 | --           |                         |      | 4           |               |     |     |     |     |     |
| 3               | NS       | Unnamed                 | --           |                         |      | 4           |               |     |     |     |     |     |
| 4               | 4        | Broadview Creek         | 4-Oct        | 205-208                 | E    | 4           |               |     |     |     |     |     |
| 5               | 5        | Piper's Creek           | 6-Oct        | -175-0                  | E    | 2           |               |     |     |     | 32  | 2   |
| 5               | 25       | Piper's Creek           | 6-Oct        | 0-42                    | E    | 2           | 1             | 3   | 13  | 10  | 4   |     |
| 5               | 6        | Piper's Creek           | 6-Oct        | 42-74                   | E    | 2           |               | 12  | 13  |     |     |     |
| 5               | 7        | Piper's Creek           | 6-Oct        | 250-300                 | E    | 2           | 6             | 71  |     |     |     |     |
| 5               | 8        | Piper's Creek           | 6-Oct        | 550-600                 | E    | 2           | 3             | 47  |     |     |     |     |
| 5               | 510      | Venema Creek            | 19-Oct       | 0-50                    | E    | 3           |               | 11  |     |     |     |     |
| 5               | 20       | Mohlendorph Creek       | 20-Oct       | 0-20                    | E    | 3           |               |     |     |     |     |     |
| 5               | 516      | Mohlendorph Creek       | 19-Oct       | 100-150                 | E    | 4           |               |     |     |     |     |     |
| 5               | 514      | Venema Creek            | 20-Oct       | 270-320                 | E    | 3           |               | 5   |     |     |     |     |
| 5               | 18       | Venema Creek            | 20-Oct       | 600-650                 | E    | 3           |               |     |     |     |     |     |
| 5               | 501      | Piper's Creek           | 19-Oct       | 820-870                 | E    | 2           |               | 35  |     |     |     |     |
| 5               | 21       | Unnamed trib H          | 11-Oct       | 0-50                    | E    | 3           |               | 3   |     |     |     |     |
| 5               | 22       | Unnamed trib H          | 11-Oct       | 250-300                 | E    | 3           |               |     |     |     |     |     |
| 5               | 10       | Piper's Creek           | 6-Oct        | 1,300-1,353             | E    | 3           |               | 55  |     |     |     |     |
| 5               | 23       | Unnamed trib K          | 11-Oct       | 0-54                    | E    | 3           |               | 5   |     |     |     |     |
| 5               | 11       | Piper's Creek           | 11-Oct       | 1,600-1,650             | E    | 3           |               | 4   |     |     |     |     |
| 5               | 12       | Piper's Creek           | 11-Oct       | 1,750-1,800             | E    | 3           |               |     |     |     |     |     |
| 5               | 24       | Unnamed trib M          | 11-Oct       | 0-7.5                   | E    | 4           |               |     |     |     |     |     |
| 5               | 13       | Piper's Creek           | 11-Oct       | 2,150-2,181             | E    | 3           |               |     |     |     |     |     |
| 5               | 14       | Piper's Creek           | 11-Oct       | 2,350-2,400             | E    | 3           |               |     |     |     |     |     |
| 6               | NS       | Unnamed                 | --           |                         |      |             |               |     |     |     |     |     |
| 7               | NS       | Unnamed                 | --           |                         |      |             |               |     |     |     |     |     |
| 8               | 32       | Blue Ridge Creek        | 25-Oct       | 350-360                 | E    | 3           |               |     |     |     |     |     |
| 8               | 33       | Blue Ridge Creek        | 25-Oct       | 1,050-1,060             | E    | 3           |               |     |     |     |     |     |
| 8               | 30       | East Fork Blue Ridge Cr | 25-Oct       | 50-70                   | E    | 3           |               |     |     |     |     |     |
| 8               | 31       | East Fork Blue Ridge Cr | 25-Oct       | 400-410                 | E    | 3           |               |     |     |     |     |     |
| 9               | 35       | Unnamed                 | 25-Oct       | 50-52                   | E    | 4           |               |     |     |     |     |     |
| 10              | 36       | Unnamed                 | 12-Oct       | 100-130                 | E    | 4           |               |     |     |     |     |     |
| 11              | NS       | Unnamed                 | --           |                         |      | 5           |               |     |     |     |     |     |
| 12              | 40       | Golden Gardens Stream B | 14-Oct       | 0                       | NF   | 4           |               |     |     |     |     |     |
| 12              | 41       | Golden Gardens Stream B | 14-Oct       | 0                       | NF   | 4           |               |     |     |     |     |     |
| 12              | 42       | Golden Gardens Stream E | 14-Oct       | 20-81                   | E    | 4           |               |     |     |     |     |     |
| 12              | 43       | Golden Gardens Stream D | 14-Oct       | 0                       | NF   | 4           |               |     |     |     |     |     |
| 12              | 44       | Golden Gardens Stream A | 14-Oct       | 0                       | NF   | 4           |               |     |     |     |     |     |
| 13              | 45       | Licton Springs          | 25-Oct       | 1,600-1,622             | E    | 3           |               |     |     |     |     |     |

TABLE 2.3.-- Number of fish collected or observed (snorkeler counts) during winter/spring surveys, February-April 2006. Appendix number is the map which displays the location of each sample site. SB = South Branch. Gear used was either electrofishing (E), beach seine (B), or snorkel survey (S). Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are cutthroat trout (CUT), rainbow trout (RBT), Chinook salmon (CHK), threespine stickleback (STB), prickly sculpin (PKS), largemouth bass (LMB), and sunfish (SUN, pumpkinseed and juvenile sunfish combined).

| Region              | Sample   |     | Stream name | Date Sampled                     | Distance from mouth (m) | Gear        | Stream Type | Number caught |     |     |     |     |     |     |     |     |
|---------------------|----------|-----|-------------|----------------------------------|-------------------------|-------------|-------------|---------------|-----|-----|-----|-----|-----|-----|-----|-----|
|                     | stream # | #   |             |                                  |                         |             |             | Appendix      | CHK | CUT | RBT | STB | CRS | PKS | LMB | SUN |
| <b>Northwest</b>    |          |     |             |                                  |                         |             |             |               |     |     |     |     |     |     |     |     |
|                     | 5        | 312 | 1.3         | Piper's Creek                    | 7-Feb                   | 1,400-1,550 | E           | 3             |     | 3   |     |     |     |     |     |     |
|                     | 5        | 311 | 1.2         | Venema Creek                     | 7-Feb                   | 600-630     | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 5        | 310 | 1.2         | Mohlendorph Creek                | 7-Feb                   | 0-35        | E           | 3             |     | 4   |     |     |     |     |     |     |
|                     | 5        | 321 | 1.2         | Mohlendorph Creek                | 7-Feb                   | 250-255     | E           | 4             |     |     |     |     |     |     |     |     |
|                     | 13       | 309 | 1.7         | Licton Springs                   | 13-Apr                  | 1,600-1,690 | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 13       | 322 | 1.7         | Licton Springs                   | 13-Apr                  | 1,800-1,830 | E           | 3             |     |     |     |     |     |     |     |     |
| <b>Northeast</b>    |          |     |             |                                  |                         |             |             |               |     |     |     |     |     |     |     |     |
|                     | 14       | 316 | 1.8         | Matthews Creek                   | 8-Feb                   | 100-115     | E           | 3             |     | 46  |     |     |     |     |     |     |
|                     | 14       | 303 | 1.11        | Littlebrook Creek                | 8-Feb                   | 350-370     | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 14       | 338 | 1.12        | Unnamed trib at rkm 5.28         | 8-Feb                   | 25-35       | E           | 4             |     |     |     |     |     |     |     |     |
|                     | 14       | 334 | 1.13        | Littles Creek                    | 8-Feb                   | 1,100-1,147 | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 14       | 335 | 1.13        | Littles Creek Pond               | 8-Feb                   | 1,200-1,225 | S           | 3             |     |     |     |     |     |     |     |     |
|                     | 14       | 323 | 1.13        | Jackson Park Golf Course Pond A  | 8-Feb                   | 6,650       | S           | 3             |     |     |     |     |     |     |     | 34  |
|                     | 14       | 336 | 1.13        | Jackson Park Golf Course Pond B  | 8-Feb                   | 6,750       | S           | 3             |     |     |     |     |     |     |     |     |
|                     | 14       | 337 | 1.13        | Jackson Park Golf Course Pond C  | 8-Feb                   | 6,850       | S           | 3             |     | 14  |     |     |     |     | 1   | 3   |
|                     | 14       | 314 | 1.10        | Meadowbrook Creek                | 8-Feb                   | 0-50        | E           | 3             |     | 2   |     |     |     |     |     |     |
|                     | 14       | 313 | 1.10        | Meadowbrook Creek Pond           | 8-Feb                   | 350-450     | B           | 3             |     |     |     |     |     |     |     |     |
|                     | 14       | 320 | 1.16        | Victory Creek                    | 8-Feb                   | 550-580     | E           | 4             |     |     |     |     |     |     |     |     |
|                     | 14       | 315 | 1.15        | Willow Creek                     | 8-Feb                   | 325-350     | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 14       | 302 | 1.16        | Park and Ride Pond (SB Thornton) | 15-Feb                  | 4,250-4,265 | E           | 3             |     | 3   |     | 1   |     |     |     |     |
| <b>East-central</b> |          |     |             |                                  |                         |             |             |               |     |     |     |     |     |     |     |     |
|                     | 27       | 316 | 1.21        | Washington Park Creek            | 15-Feb                  | 0-25        | E           | 3             |     |     |     |     |     |     |     | 1   |
|                     | 27       | 324 | 1.21        | Washington Park Creek            | 15-Feb                  | 100-105     | E           | 3             |     | 1   |     |     |     |     |     |     |
|                     | 27       | 325 | 1.21        | Washington Park Creek            | 15-Feb                  | 850-875     | E           | 3             |     |     |     |     |     |     |     |     |
| <b>Southwest</b>    |          |     |             |                                  |                         |             |             |               |     |     |     |     |     |     |     |     |
|                     | 34       | 333 | 1.24        | Longfellow Creek                 | 15-Feb                  | 1,000-1,028 | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 34       | 317 | 1.24        | Longfellow Creek                 | 15-Feb                  | 1,650-1,718 | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 35       | 319 | 1.25        | Puget Creek                      | 9-Feb                   | 0-60        | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 35       | 318 | 1.25        | Unnamed trib at rkm 0.2          | 9-Feb                   | 3-6.5       | E           | 3             |     |     | 1   |     |     |     |     |     |
|                     | 44       | 326 | 1.28        | Durham Creek                     | 9-Feb                   | 450-487     | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 45       | 327 | 1.28        | Durham Creek B                   | 9-Feb                   | 0-10        | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 46       | 328 | 1.28        | Durham Creek C                   | 9-Feb                   | 0-50        | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 43       | 329 | 1.28        | Seola Beach Creek                | 9-Feb                   | 175-185     | E           | 4             |     |     |     |     |     |     |     |     |
| <b>Southeast</b>    |          |     |             |                                  |                         |             |             |               |     |     |     |     |     |     |     |     |
|                     | 48       | 330 | 1.29        | Mapes Creek (Kabota Gardens)     | 9-Feb                   | 1,480-1,680 | E,B         | 3             |     |     |     |     |     |     |     |     |
|                     | 49       | 300 | 1.30        | Taylor Creek                     | 16-Feb                  | 1,060-1,110 | E           | 3             |     |     |     |     |     |     |     |     |
|                     | 49       | 307 | 1.30        | Taylor Creek                     | 13-Apr                  | 0-34        | E           | 2             | 1   | 6   |     | 2   | 28  | 3   |     |     |
|                     | 49       | 308 | 1.30        | Taylor Creek                     | 13-Apr                  | 34-64       | E           | 3             |     | 6   |     |     |     |     |     |     |

Northeast region. -- In Thornton Creek, we captured fish from the mouth up to the headwaters of the north and south branches (Tables 2.4 and 2.5). Over 52% of the fish we collected in mainstem habitats of the Thornton Creek system were cutthroat trout, which ranged throughout the system. Other native species collected in Thornton Creek were coho salmon, rainbow trout, threespine stickleback, lamprey, prickly sculpin, and coastrange sculpin. We also captured several introduced fish species including rock bass, largemouth bass, and pumpkinseed. The only introduced fish collected in the South Branch of Thornton Creek system were some pumpkinseed, which were only captured in Kramer Creek. We also captured fish in six different tributaries of Thornton Creek (Matthews, Maple, Meadowbrook Park, Kramer, Willow, and Littlebrook Creeks). Fish captured in the tributaries were also mainly cutthroat trout. Fish distributions in the tributaries appeared to be limited by barriers, such as culverts and other man-made obstacles. Most tributary fish were captured within 100 m of the mouth. In Matthews Creek and Maple Creek, we only captured fish downstream of the culvert at Sand Point Way to the confluence with Thornton Creek. On Littlebrook Creek, we only captured fish in a small section downstream from the culvert under 115<sup>th</sup> St to the confluence with the North Branch of Thornton Creek. In Willow Creek, we only captured fish downstream of the culvert under 98<sup>th</sup> St. to the confluence with the South Branch of Thornton Creek.

During the fish distribution surveys, 15 coho salmon were collected between the mouth of Thornton Creek to the Meadowbrook Retention Pond intake structure at Rm 2,000. Upstream only one coho salmon was ever collected, which was captured during our reference site sampling on Kramer Creek (200 m upstream from the Meadowbrook Retention Pond intake structure).

Ravenna Park Creek was the only other creek in the northeast region in which we collected any fish. We collected two rainbow trout that were approximately 200 m upstream from where the creek enters the culvert at the south end of Ravenna Park.

Yesler Creek (Stream #16) was not sampled due to inaccessibility and Inverness Creek (Stream #15) was sampled but no fish were captured.

TABLE 2.4.-- Number of fish collected during fish distribution surveys in the Thornton Creek (stream system #14) mainstem (Rm 0 – 2,200) and the North Branch of Thornton Creek (Rm 2,200-6,650), August-October 2005. Gear used was either electrofishing (E) or beach seine (B). Locations of sample sites are displayed in Appendices 1.8-1.13. Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are coho salmon (COH), cutthroat trout (CUT), threespine stickleback (STB), bass (rock bass and largemouth bass combined; BAS), sunfish (SUN), pumpkinseed and juvenile sunfish combined), coastrange sculpin (CRS), prickly sculpin (PKS), lamprey ammocoetes (LPU), and oriental weatherfish (OWF).

| Sample # | Stream name                     | Date Sampled | Distance from mouth (m) | Gear | Stream Type | Number caught |     |     |     |     |     |     |     |     |  |
|----------|---------------------------------|--------------|-------------------------|------|-------------|---------------|-----|-----|-----|-----|-----|-----|-----|-----|--|
|          |                                 |              |                         |      |             | COH           | CUT | STB | BAS | SUN | CRS | PKS | LPU | OWF |  |
| 50       | Thornton Creek - mainstem       | 22-Aug       | 30-80                   | E,B  | 2           | 6             |     | 3   |     |     |     |     | 74  |     |  |
| 68       | Matthews Creek Pond             | 22-Aug       | 15-37                   | B    | 3           |               |     | 58  |     |     |     |     | 2   |     |  |
| 268      | Matthews Creek                  | 22-Aug       | 37-52                   | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 70       | Unnamed trib C at rkm 0.54      | 29-Aug       | 0-30                    | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 51       | Thornton Creek - mainstem       | 22-Aug       | 250-300                 | E    | 2           |               | 7   |     |     |     | 74  | 46  |     |     |  |
| 69       | Maple Creek                     | 22-Aug       | 0-50                    | E    | 3           |               | 13  |     |     |     |     |     |     |     |  |
| 71       | Unnamed trib A at rkm 0.09      | 29-Aug       | 340-390                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 73       | Unnamed trib A at rkm 0.09      | 29-Aug       | 740-790                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 74       | Unnamed trib B at rkm 0.08      | 29-Aug       | 360-410                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 72       | Maple Creek                     | 29-Aug       | 450-500                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 75       | Maple Creek                     | 20-Sep       | 850-890                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 52       | Thornton Creek - mainstem       | 31-Aug       | 350-450                 | E    | 2           | 1             | 25  |     | 1   |     | 201 | 111 | 4   | 2   |  |
| 53       | Thornton Creek - mainstem       | 30-Aug       | 500-600                 | E    | 2           | 9             | 135 | 43  |     | 5   | 7   |     |     |     |  |
| 54       | Thornton Creek - mainstem       | 12-Sep       | 1,100-1,150             | E    | 2           | 1             | 67  | 31  |     | 17  | 1   |     |     |     |  |
| 76       | Mock Creek                      | 15-Sep       | 460-480                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 77       | Mock Creek                      | 15-Sep       | 770-800                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 55       | Thornton Creek - mainstem       | 15-Aug       | 2,000-2,045             | E    | 2           | 7             | 421 | 1   | 3   | 2   | 1   |     |     |     |  |
| 56       | North Branch Thornton           | 12-Sep       | 2,850-2,901             | E    | 2           |               | 142 |     |     | 2   |     |     |     |     |  |
| 78       | Littlebrook Creek               | 8-Sep        | 0-50                    | E    | 3           |               | 26  |     |     |     |     |     |     |     |  |
| 79       | Littlebrook Creek               | 8-Sep        | 350-400                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 80       | Littlebrook Creek               | 12-Sep       | 800-820                 | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 81       | Littlebrook Creek               | 23-Sep       | 2,350-2,375             | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 82       | Littlebrook Creek               | 23-Sep       | 2,930-2,970             | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 83       | Littles Creek                   | 5-Oct        | 1,100-1,147             | E    | 3           |               |     |     |     |     |     |     |     |     |  |
| 84       | Littles Creek Pond              | 5-Oct        | 1,200-1,225             | B    | 3           |               |     |     |     |     |     |     |     |     |  |
| 57       | North Branch Thornton           | 17-Sep       | 3,650-3,700             | E    | 2           |               | 111 |     |     | 4   |     |     |     |     |  |
| 58       | North Branch Thornton           | 19-Sep       | 4,300-4,363             | E    | 2           |               | 80  |     |     | 2   |     |     |     |     |  |
| 59       | North Branch Thornton           | 7-Sep        | 5,150-5,200             | E    | 2           |               | 42  |     |     |     |     |     |     |     |  |
| 60       | North Branch Thornton           | 7-Sep        | 5,950-5,970             | E    | 2           |               | 15  |     |     | 1   |     |     |     |     |  |
| 61       | North Branch Thornton           | 7-Sep        | 6,600-6,650             | E    | 2           |               | 125 |     | 1   | 27  |     |     |     |     |  |
| 85       | Jackson Park Golf Course Pond A | 26-Sep       | 6,650                   | E,B  | 3           |               |     |     |     | 73  |     |     |     |     |  |
| 285      | Jackson Park Golf Course Pond B | 26-Sep       | 6,750                   | E,B  | 3           |               |     |     |     |     |     |     |     |     |  |
| 286      | Jackson Park Golf Course Pond C | 26-Sep       | 6,850                   | E,B  | 3           |               |     |     | 6   | 6   |     |     |     |     |  |

TABLE 2.5.-- Number of fish collected during fish distribution surveys in the South Branch of Thornton Creek and other streams (stream system #'s 15-17) in the northeast region, September-October 2005. NS = no sample. NSCC = North Seattle Community College. Gear used was either electrofishing (E), beach seine (B), or gill nets (G). Locations of sample sites are displayed in Appendices 1.10, 1.14-1.18. Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are coho salmon (COH), cutthroat trout (CUT), rainbow trout (RBT), and threespine stickleback (STB).

| Stream system # | Sample # | Stream name              | Date Sampled | Distance from mouth (m) | Gear | Stream Type | Number caught |     |     |     |
|-----------------|----------|--------------------------|--------------|-------------------------|------|-------------|---------------|-----|-----|-----|
|                 |          |                          |              |                         |      |             | COH           | CUT | RBT | STB |
| 14              | 62       | South Branch Thornton    | 27-Sep       | 0-45                    | E    | 2           | 51            |     |     |     |
| 14              | 86       | Meadowbrook Creek        | 20-Sep       | 0-56                    | E    | 3           |               |     |     |     |
| 14              | 88       | Meadowbrook Pond         | 20-Sep       | 350-500                 | B, E | 3           |               |     |     |     |
| 14              | 524      | Kramer Creek             | 28-Sep       | 0-108                   | E    | 3           | 1             | 35  |     |     |
| 14              | 521      | South Branch Thornton    | 13-Oct       | 550-688                 | E    | 2           | 177           |     |     |     |
| 14              | 89       | Unnamed trib C           | 27-Sep       | 0-6.8                   | E    | 4           |               |     |     |     |
| 14              | 90       | Willow Creek             | 20-Sep       | 0-50.5                  | E    | 3           | 38            |     |     |     |
| 14              | 91       | Willow Creek             | 27-Sep       | 250-265                 | E    | 3           | 29            |     |     |     |
| 14              | 95       | Unnamed trib E           | 21-Sep       | 0-32                    | E    | 3           | 12            |     |     |     |
| 14              | 96       | Unnamed trib E           | 21-Sep       | 350-385.5               | E    | 3           |               |     |     |     |
| 14              | 92       | Willow Creek             | 21-Sep       | 350-388                 | E    | 3           |               |     |     |     |
| 14              | 93       | Willow Creek             | 21-Sep       | 920-972                 | E    | 3           |               |     |     |     |
| 14              | 94       | Willow Creek             | 21-Sep       | 1,500-1,518             | E    | 3           |               |     |     |     |
| 14              | 64       | South Branch Thornton    | 21-Sep       | 1,350-1,410             | E    | 2           | 95            |     |     |     |
| 14              | 65       | South Branch Thornton    | 22-Sep       | 2,250-2,297             | E    | 2           | 29            |     |     | 13  |
| 14              | 97       | Victory Creek            | 27-Sep       | 250-283                 | E    | 3           |               |     |     |     |
| 14              | 66       | South Branch Thornton    | 22-Sep       | 3,200-3,247             | E    | 2           | 17            |     | 1   | 4   |
| 14              | 67       | South Branch Thornton    | 22-Sep       | 3,750-3,807             | E    | 2           | 1             |     | 78  |     |
| 14              | 98       | Park and Ride Pond North | 26-Oct       | 4,250-4,270             | B, G | 3           | 180           |     |     |     |
| 14              | 99       | NSCC Pond                | 1-Sep        | 4,600-4,620             | B, G | 3           | 161           |     |     |     |
| 14              | 199      | NSCC Slough              | 1-Sep        | 4,650-4,670             | B, E | 3           | 47            |     |     |     |
| 15              | 100      | Iverness Creek           | 28-Sep       | 0-35                    | E    | 4           |               |     |     |     |
| 16              | NS       | Yesler Creek             | --           |                         |      | 3           |               |     |     |     |
| 17              | 105      | Ravenna Park Creek       | 4-Oct        | 175-225                 | E    | 3           |               |     |     | 2   |
| 17              | 106      | Ravenna Park Creek       | 5-Oct        | 600-650                 | E    | 3           |               |     |     |     |



*West-central region.*-- The west-central region contained nine streams (Streams #'s 18–26), of which only four locations in Discovery Park were found to have any fish (Table 2.6). Three of the locations were slow-water habitats of the Scheuerman Creek drainage, and several threespine stickleback were captured at each location. The first site was a man-made pool (14-m long by 5-m wide; maximum depth 0.34) at Rm 100. The other two locations were small ponds (maximum depths, 1.5 and 1.0 m) on a left-bank tributary. A third pond at the headwaters of this tributary did not contain any fish. For the three locations combined, a total of 113 threespine stickleback were captured with a beach seine. We also collected a single goldfish in another pond located in the center of Discovery Park. We were unable to survey Streams #'s 23-25 due to inaccessibility. We did not capture any fish in Streams #'s 18–20, 22, and 26.

TABLE 2.6.-- Number of fish collected in the west-central region watersheds during fish distribution surveys, August-October 2005. NS = no sample. Gear used was either electrofishing (E), beach seine (B), or not fished (NF, only habitat information collected). Locations of sample sites are displayed in Appendices 1.19 and 1.20. Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are goldfish (GDF) and threespine stickleback (STB).

| Stream system # | Sample # | Stream name                      | Date Sampled | Distance from mouth (m) | Gear | Stream Type | Number caught |     |
|-----------------|----------|----------------------------------|--------------|-------------------------|------|-------------|---------------|-----|
|                 |          |                                  |              |                         |      |             | GDF           | STB |
| 18              | 110      | Mahteen Creek                    | 19-Aug       | 0-40                    | E    | 4           |               |     |
| 19              | 111      | Lawton Creek                     | 17-Aug       | 25-75                   | E    | 4           |               |     |
| 20              | 112      | Wolfe Creek                      | 19-Aug       | 0-50                    | E    | 3           |               |     |
| 20              | 113      | Wolfe Creek                      | 24-Aug       | 400-410                 | E    | 3           |               |     |
| 21              | 114      | Scheuerman Creek                 | 25-Aug       | 0-50                    | E    | 3           |               |     |
| 21              | 115      | Scheuerman Creek                 | 25-Aug       | 130-144                 | B, E | 3           |               | 43  |
| 21              | 116      | Scheuerman Creek                 | 19-Aug       | 280-325                 | E    | 3           |               |     |
| 21              | 117      | Scheuerman Creek                 | 19-Aug       | 460-490                 | E    | 3           |               |     |
| 21              | 217      | Scheuerman Creek - pond          | 25-Aug       | 760-791                 | B    | 4           |               |     |
| 21              | 118      | Unnamed trib A at rkm 0.28       | 19-Aug       | 0-50                    | E    | 3           |               |     |
| 21              | 119      | Unnamed trib A at rkm 0.28 -pond | 25-Aug       | 200                     | B, E | 3           |               | 70  |
| 21              | 120      | Unnamed trib A at rkm 0.28 -pond | 25-Aug       | 320                     | B, E | 3           |               | 14  |
| 21              | 121      | Unnamed trib A at rkm 0.28 -pond | 25-Aug       | 400                     | E    | 3           |               |     |
| 22              | 122      | Owl's Creek                      | 29-Sep       | 0-12                    | E    | 4           |               |     |
| 22              | 123      | South Owl's Creek                | 29-Sep       | 0                       | NF   | 4           |               |     |
| 22              | 124      | Sewer Plant Beach Pond           | 3-Oct        | 0-18                    | B    | 3           |               |     |
| 22              | 125      | Unnamed Trading Post Pond        | 3-Oct        | 0                       | B    | 3           | 1             |     |
| 23              | NS       | Unnamed                          | --           |                         |      |             |               |     |
| 24              | NS       | Unnamed                          | --           |                         |      |             |               |     |
| 25              | NS       | Unnamed                          | --           |                         |      |             |               |     |
| 26              | 126      | Magnolia Blvd Stream             | 25-Aug       | 200-220                 |      | 4           |               |     |

*East-central region.*-- The east-central region contains seven streams (stream systems #'s 27-33) of which only the Washington Park Creek drainage (stream system #27) contained fish (both native and introduced species; Table 2.7). Within the stream habitat of Washington Park Creek, we only collected fish in the lower 110 m. A perched culvert at the upstream end of this section appeared to be a barrier to fish moving upstream. Cutthroat trout and prickly sculpin were collected in the plunge pool below the culvert (maximum depth 0.47 m). Threespine stickleback, prickly sculpin, and smallmouth bass were collected near the mouth of the creek. Upstream, fish were only collected in two ponds. All fish were most likely introduced into the ponds. Fourteen goldfish and seven brown bullhead were collected in a small pond on a small right-bank tributary to Washington Park Creek at Rm 1,475. Also, 37 koi (ornamental common carp) and 22 goldfish were collected in Japanese Gardens Pond at the headwaters of Washington Park Creek.

TABLE 2.7.-- Number of fish collected in the east-central region watersheds during fish distribution surveys, July-August 2005. Gear used was either electrofishing (E) or beach seine (B). Locations of sample sites are displayed in Appendices 1.21 and 1.22. Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are goldfish (GDF), common carp (koi, CRP), brown bullhead (BBH), threespine stickleback (STB), smallmouth bass (SMB), and prickly sculpin (PKS). Sample #133 is known as the Japanese Garden Pond.

| Stream system # | Sample # | Stream name                  | Date Sampled | Distance from mouth (m) | Gear | Stream Type | Number caught |     |     |     |     |
|-----------------|----------|------------------------------|--------------|-------------------------|------|-------------|---------------|-----|-----|-----|-----|
|                 |          |                              |              |                         |      |             | GDF           | CRP | BBH | STB | SMB |
| 27              | 130      | Washington Park Creek        | 11-Aug       | 0-50                    | E    | 3           |               |     | 4   | 1   | 4   |
| 27              | 230      | Washington Park Creek        | 11-Aug       | 50-105                  | E    | 3           |               |     |     |     | 3   |
| 27              | 131      | Washington Park Creek        | 11-Aug       | 850-900                 | E    | 3           |               |     |     |     |     |
| 27              | 132      | Unnamed trib A - pond        | 11-Aug       | 50-75                   | B    | 3           | 14            |     | 7   |     |     |
| 27              | 133      | Washington Park Creek - pond | 22-Aug       | 1,680                   | B    | 3           | 22            | 37  |     |     |     |
| 28              | 134      | Interlaken Creek - reach A   | 17-Aug       | 80-90                   | E    | 4           |               |     |     |     |     |
| 28              | 135      | Interlaken Creek - reach B   | 19-Aug       | 0-20                    | E    | 4           |               |     |     |     |     |
| 29              | 136      | Interlaken Creek - reach C   | 19-Aug       | 0-50                    | E    | 4           |               |     |     |     |     |
| 30              | 137      | Boren Park Area              | 19-Aug       | 0-30                    | E    | 4           |               |     |     |     |     |
| 31              | 138      | Madrona Park North Trib      | 10-Aug       | 275-295                 | E    | 4           |               |     |     |     |     |
| 32              | 139      | Madrona Park South Trib      | 10-Aug       | 50-60                   | E    | 4           |               |     |     |     |     |
| 33              | 140      | Frink Creek                  | 28-Jul       | 575-625                 | E    | 3           |               |     |     |     |     |
| 33              | 141      | Frink Creek                  | 28-Jul       | 800-820                 | E    | 3           |               |     |     |     |     |

Southwest region.-- The southwest region contains 13 streams (stream system #'s 34-46) of which Longfellow Creek, Puget Creek, Fauntleroy Creek, and Durham Creek contained fish. In Longfellow Creek, we only captured fish in a few sections (Table 2.8). At the mouth (just upstream from where Longfellow Creek enters the lowest culvert to the Duwamish River) we captured 17 threespine stickleback, 2 prickly sculpin, and 1 Pacific staghorn sculpin. We didn't catch another fish until Rm 2,100 where three unidentified trout (most likely rainbow trout) were collected. Between Rm 2,100 and 3,500, we captured three rainbow trout, and three juvenile coho salmon. No other fish were captured upstream of this point on Longfellow Creek despite adequate habitat.

Puget Creek was sampled in the summer of 2005 and no fish were collected. However in the winter surveys of 2006, we captured a 224 mm FL rainbow trout in a small pool of a tributary to Puget Creek (Figure 2.1). This tributary is 50 m upstream from the lower end of the daylighted section of Puget Creek. The tributary appears to have more streamflow than Puget Creek during base flow conditions. In the summer of 2006, we sampled this pool again but did not capture any fish.

Fauntleroy Creek is a major creek in this region that has an annual return of coho salmon. In the intertidal area, we only collected three Pacific sand lance. Upstream at the upper edge of the intertidal area, we collected four juvenile coho salmon and four Pacific staghorn sculpin. In our surveys further upstream, we only captured juvenile coho salmon, which were collected as far upstream as Rm 775 (Table 2.8).

The Durham Creek system (stream system #'s 44 – 46) was also sampled. The lowest daylighted section (stream system # 44) is located in the South Park Pea Patch where we collected 14 threespine stickleback and 3 juvenile coho salmon. This stream section has a low gradient and has adequate pool habitat and cover (large amounts of water cress *Nasturtium officinale*). Just upstream of this section is a steep culvert and no fish were observed in the stream section above the culvert. It is not clear if stream system #'s 45 and 46 are directly connected to stream system # 44. Stream system #45 is likely connected to Durham Creek but stream system #46 may be part of the Hamm Creek system. Although these streams had adequate habitat and streamflow, no fish were collected. Most likely, these streams are perched above impassable culverts.

TABLE 2.8.-- Number of fish collected in the southwest region watersheds during fish distribution surveys, August-September 2006. NS = no sample. Gear used was either electrofishing (E) or not fished (NF, only habitat information collected). Locations of sample sites are displayed in Appendices 1.23-1.28. Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are coho salmon (COH), rainbow trout (RBT), threespine stickleback (STB), prickly sculpin (PKS), and Pacific staghorn sculpin (SHS).

| Stream system # | Sample # | Stream name             | Date Sampled | Distance from mouth (m) | Gear | Stream Typing | Number caught |     |     |     |     |
|-----------------|----------|-------------------------|--------------|-------------------------|------|---------------|---------------|-----|-----|-----|-----|
|                 |          |                         |              |                         |      |               | COH           | RBT | STB | PKS | SHS |
| 34              | 150      | Longfellow Creek        | 1-Aug        | 0-80                    | E    | 3             |               |     | 17  | 2   | 1   |
| 34              | 146      | Golf Course trib        | 1-Aug        | 0-27                    | E    | 4             |               |     |     |     |     |
| 34              | 147      | Unnamed trib A          | 1-Aug        | 225-245                 | E    | 4             |               |     |     |     |     |
| 34              | 151      | Longfellow Creek        | 1-Aug        | 1,000-1,050             | E    | 3             |               |     |     |     |     |
| 34              | 152      | Longfellow Creek        | 1-Aug        | 1,650-1,685             | E    | 3             |               |     |     |     |     |
| 34              | 153      | Longfellow Creek        | 2-Aug        | 2,150-2,220             | E    | 3             |               |     |     |     |     |
| 34              | 154      | Longfellow Creek        | 2-Aug        | 2,850-2,917             | E    | 3             | 3             | 2   |     |     |     |
| 34              | 155      | Longfellow Creek        | 2-Aug        | 3,400-3,450             | E    | 3             |               | 1   |     |     |     |
| 34              | 156      | Longfellow Creek        | 2-Aug        | 3,775-3,822             | E    | 3             |               |     |     |     |     |
| 34              | 148      | Unnamed trib B          | 2-Aug        | 0-10                    | E    | 5             |               |     |     |     |     |
| 34              | 149      | Unnamed trib C          | 3-Aug        | 0-3                     | E    | 4             |               |     |     |     |     |
| 34              | 257      | Longfellow Creek        | 3-Aug        | 4,175-4,225             | E    | 3             |               |     |     |     |     |
| 34              | 157      | Longfellow Creek        | 3-Aug        | 4,850-4,898             | E    | 3             |               |     |     |     |     |
| 34              | 158      | Longfellow Creek        | 3-Aug        | 5,300-5,347             | E    | 3             |               |     |     |     |     |
| 34              | 159      | Longfellow Creek        | 4-Aug        | 5,600-5,647             | E    | 3             |               |     |     |     |     |
| 35              | 160      | Puget Creek             | 9-Aug        | 0-54                    | E    | 3             |               |     |     |     |     |
| 35              | 161      | Unnamed trib A at rm 12 | 9-Aug        | 0-6.5                   | E    | 4             |               |     |     |     |     |
| 35              | 162      | Unnamed trib B at rm 54 | 9-Aug        | 0-13.5                  | E    | 3             |               |     |     |     |     |
| 36              | 163      | Unnamed Creek           | 28-Jul       |                         | NF   | 5             |               |     |     |     |     |
| 37              | 164      | Fairmount Creek         | 28-Jul       | 275-280                 | E    | 3             |               |     |     |     |     |
| 38              | 165      | Schmitz Creek           | 10-Aug       | 0-50                    | E    | 3             |               |     |     |     |     |
| 38              | 166      | Schmitz Creek           | 10-Aug       | 250-300                 | E    | 3             |               |     |     |     |     |
| 38              | 167      | Schmitz Creek           | 10-Aug       | 575-625                 | E    | 3             |               |     |     |     |     |
| 38              | 170      | Unnamed trib G          | 10-Aug       | 0-15                    | E    | 4             |               |     |     |     |     |
| 38              | 168      | Schmitz Creek           | 10-Aug       | 675-710                 | E    | 3             |               |     |     |     |     |
| 38              | 171      | Unnamed trib K          | 10-Aug       | 85-100                  | E    | 4             |               |     |     |     |     |
| 38              | 169      | Schmitz Creek           | 10-Aug       | 710-725                 | E    | 4             |               |     |     |     |     |
| 39              | 172      | Mee-Kwa-Mooks Creek     | 10-Aug       | 130-140                 | E    | 4             |               |     |     |     |     |
| 39              | 272      | Unnamed trib A          | 10-Aug       | 0-10                    | E    | 4             |               |     |     |     |     |
| 40              | 173      | Pelly Creek             | 10-Aug       | 130-140                 | E    | 4             |               |     |     |     |     |
| 41              | NS       | Unnamed Stream          | --           |                         | E    |               |               |     |     |     |     |
| 42              | 180      | Fauntleroy Creek        | 18-Aug       | 0-85                    | E    | 3             | 4             |     |     |     | 4   |
| 42              | 181      | Fauntleroy Creek        | 18-Oct       | 250-353                 | E    | 3             | 10            |     |     |     |     |
| 42              | 182      | Fauntleroy Creek        | 23-Aug       | 450-475                 | E    | 3             | 4             |     |     |     |     |
| 42              | 183      | Fauntleroy Creek        | 18-Aug       | 725-775                 | E    | 3             | 9             |     |     |     |     |
| 42              | 184      | Unnamed trib A          | 18-Aug       | 0-20                    | E    | 4             |               |     |     |     |     |
| 43              | NS       | Seola Beach Creek (Dry) | --           |                         | E    | 5             |               |     |     |     |     |
| 44              | 190      | Durham Creek            | 9-Aug        | 80-100                  | E    | 3             | 4             |     | 12  |     |     |
| 44              | 191      | Durham Creek            | 9-Aug        | 150-200                 | E    | 3             |               |     |     |     |     |
| 45              | 193      | Durham Creek            | 2-Aug        | 450-487                 | E    | 3             |               |     |     |     |     |
| 45              | 194      | Durham Creek B          | 2-Aug        | 0-10                    | E    | 3             |               |     |     |     |     |
| 46              | 192      | Durham Creek C          | 2-Aug        | 0-50                    | E    | 3             |               |     |     |     |     |



FIGURE 2.1.—Photo of the rainbow trout captured in a small tributary of Puget Creek during the winter surveys of 2006.

*Southeast region.*-- This region only contains three streams (stream systems #'s 47-49): Mt. Baker Creek, Mapes Creek, and Taylor Creek. No fish were collected in Mt. Baker Creek. At Mapes Creek, we sampled at Rm 1,100 and the Kubota Garden Ponds at Rm 1,500. The only fish collected in Mapes Creek were two threespine stickleback collected at Rm 1,100.

Taylor Creek is the largest creek in this section and flows north into Lake Washington. We sampled from the mouth of the creek at Lake Washington to the headwaters of the west fork (Table 2.9). In the lower 50 m of the stream, we collected one juvenile Chinook salmon, 20 juvenile coho salmon, 29 coastrange sculpin, 30 cutthroat trout, and 17 threespine stickleback. We also observed a single unidentified lamprey just above our site. Upstream from this site, the only fish we collected were juvenile coho salmon and cutthroat trout (Table 2.9). We did not collect any fish in the east fork of Taylor Creek, but we captured three cutthroat trout in the west fork in a small pool close to the confluence. This was the furthest upstream site where we captured fish on Taylor Creek.

TABLE 2.9.-- Number of fish collected in the southeast region watersheds during fish distribution surveys, August-September 2006. Gear used was either electrofishing (E) or not fished (NF, only habitat information collected). Locations of sample sites are displayed in Appendices 1.29-1.30. Stream type indicates the Washington State Department of Natural Resources interim water typing classification system on a scale of 1 to 5. Fish species (number caught) are Chinook salmon (CHK), coho salmon (COH), cutthroat trout (CUT), threespine stickleback (STB), and coastrange sculpin (CRS).

| Stream system # | Sample # | Stream name             | Date Sampled | Distance from mouth (m) | Gear | Stream Type | Number caught |     |     |     |     |
|-----------------|----------|-------------------------|--------------|-------------------------|------|-------------|---------------|-----|-----|-----|-----|
|                 |          |                         |              |                         |      |             | CHK           | COH | CUT | STB | CRS |
| 47              | 200      | Mt. Baker Creek         | 25-Jul       | 195-280                 | E    | 4           |               |     |     |     |     |
| 48              | 201      | Mapes Creek             | 25-Jul       | 900-935                 | E    | 3           |               |     |     |     | 2   |
| 48              | 202      | Mapes Creek             | 26-Jul       | 1,415-1,515             | E    | 3           |               |     |     |     |     |
| 48              | 209      | Unnamed trib at rkm 1.6 | 26-Jul       | 50-75                   | E    | 4           |               |     |     |     |     |
| 49              | 203      | Taylor Creek            | 30-Jun       | 0-50                    | E    | 2           | 1             | 20  | 30  | 17  | 33  |
| 49              | 204      | Taylor Creek            | 30-Jun       | 415-465                 | E    | 2           |               | 51  | 9   |     |     |
| 49              | 205      | Taylor Creek            | 30-Jun       | 825-875                 | E    | 3           |               | 1   | 5   |     |     |
| 49              | 206      | East Fork Taylor Creek  | 7-Jul        | 0-50                    | E    | 3           |               |     |     |     |     |
| 49              | 207      | East Fork Taylor Creek  | 19-Jul       | 50-100                  | E    | 3           |               |     |     |     |     |
| 49              | 208      | West Fork Taylor Creek  | 19-Jul       | 0-50                    | E    | 3           |               |     |     | 3   |     |
| 49              | 210      | West Fork Taylor Creek  | 19-Jul       | 1,150                   | NF   | 3           |               |     |     |     |     |

### *Fish Distribution by Species*

*Petromyzontidae, lampreys.*—During fish distribution surveys, we only collected lamprey (ammocoetes) in Thornton Creek at Rm 350-400. We did observe one unidentified adult lamprey in Taylor Creek (Rm 0.7) during winter surveys but it was not captured. Additional surveys for lamprey were conducted in the lower reach of Taylor Creek and Fautleroy Creek in September 2006 with a lamprey electrofishing unit; however, no lamprey was collected. Further surveys with the lamprey electrofishing unit are needed; however, initial results indicate lampreys are rare in Seattle streams. Species identification of lamprey was not done but they were most likely western brook lamprey, which have been observed in other Lake Washington basin streams.

Salmonidae, salmon and trout.-- Cutthroat trout was the dominant salmonid species captured during the ichthyofauna surveys. They were common in Thornton Creek, Taylor Creek, and Piper's Creek drainages. Additionally, a large cutthroat trout (211 mm FL) was collected in the lower section on Longfellow Creek (reference site; see Chapter 3) and another cutthroat trout (103 mm FL) was collected near the mouth of Washington Park Creek during 2006 winter surveys (Table 2.3). In the Thornton Creek drainage they were present from the mouth upstream to the headwaters of both the North and South Forks, including the tributaries Matthews Creek, Maple Creek, Meadowbrook Park Creek, Willow Creek, and Littlebrook Creek (Figure 2.2). Three cutthroat trout were also collected from a small pond by the Park and Ride east of I-5, near 1<sup>st</sup> Ave and 100<sup>th</sup> St. during the 2006 winter surveys (Table 2.3; Figure 2.3). We also documented the presence of cutthroat trout in Jackson Park Golf Course pond C during winter surveys (Table 2.3). Overall, cutthroat trout collected made up over 53% of the total number of fish caught throughout the Thornton Creek drainage ( $n = 1,481$ , mean FL = 93.3 mm; range, 40-291 mm FL) and was the dominant species caught at most sites (Figure 2.4). In the Piper's Creek drainage, they were found from the mouth to Rm 1,850 on the mainstem, in Venema Creek to Rm 325, and in the lower 50 m of Mohlendorph Creek (Figure 2.2). Cutthroat trout were also collected from two other tributaries of Piper's Creek located at Rm 975 and Rm 1,450. Cutthroat trout made up over 71% of the total number of fish captured in the Piper's Creek drainage ( $n = 207$ ; mean FL, 97.2 mm; range, 55-215 mm FL; Figure 2.5). Cutthroat trout were also collected in the Taylor Creek drainage from the mouth up to Rm 1,000 and at the mouth of the West Fork at Rm 1,100. Cutthroat trout represented 27% of the fish collected in the Taylor Creek drainage ( $n = 58$ ; mean FL, 114.6 mm; range 47-235 mm FL).




FIGURE 2.2.—Distribution of cutthroat trout in Seattle's streams, 2005-2006.





FIGURE 2.3.— Photo of cutthroat trout captured in a small pond by the Park and Ride just east of I-5, near 1<sup>st</sup> Ave and 100<sup>th</sup> St. during the 2006 winter surveys.




FIGURE 2.4.-- Relative abundance of the three main types of fish in the Thornton Creek system, August-October 2005. The distance from mouth (m) is the distance from the stream mouth on Lake Washington. In some cases, the total percent for a site may not add to 100% because other fish (lamprey, coho salmon, and centrarchids) are not displayed. At Rm 2,200, the South Branch and the North Branch merge together to form the lower mainstem of Thornton Creek. The North Branch is not displayed; cutthroat trout was the dominant fish species at each site (mean, 95% of catch; range 82-100%) in the North Branch.




FIGURE 2.5. Relative abundance of the three main types of fish in Piper’s Creek (mainstem only), October 2005. The distance from mouth (m) is the distance from the stream mouth on Puget Sound.

Few rainbow trout were captured in both the summer and winter sampling. Small numbers of rainbow trout were found in Thornton Creek, Ravenna Creek, and Longfellow Creek during the 2005 summer surveys (Figure 2.6). Additionally, a single rainbow trout was collected in Puget Creek during the winter surveys. Rainbow trout captured in Thornton Creek, Longfellow Creek, and perhaps other locations seemed to have some phenotypic characteristics of cutthroat trout (i.e., somewhat longer bottom jaw and very faint “slash” marks under lower jaw) and may have been hybrids. Genetic analysis of these individuals is needed to help determine their identification.



FIGURE 2.6.— Photos of rainbow trout captured in Longfellow Creek (top photo) and Ravenna Creek (bottom photo) during the 2005 summer surveys.

Juvenile coho salmon were captured in Piper's Creek, Thornton Creek, Kramer Creek, Longfellow Creek, Fauntleroy Creek, Taylor Creek, and a section of Durham Creek (Figure 2.7). In Piper's Creek they were present from the mouth to Rm 625. Coho salmon only made up 3.4% of the total number of fish collected in the Piper's Creek drainage ( $n = 10$ ; mean FL, 81.1 mm, range, 78-88 mm FL). Coho salmon were also collected in the Thornton Creek drainage, but were only found in the lower 2 km of the mainstem and two individuals were collected from Kramer Creek during our depletion-removal sampling. Less than 1% of the total catch in the Thornton Creek drainage was coho salmon ( $n = 24$ ; mean FL, 84 mm; range, 72-97 mm FL). During fish distribution surveys, only two coho salmon were captured in Longfellow Creek at Rm 2,850 and four in the Durham Creek (Table 2.8). In the Fauntleroy Creek drainage, coho salmon were captured up to Rm 775 (Table 2.8). There is a barrier at Rm 400 under California Ave SW, and coho salmon found above this point were most likely stocked there by local enhancement groups. Coho were the dominant species present and made up 73% of all fish collected in the Fauntleroy Creek drainage ( $n = 19$ ; mean FL, 68.2 mm; range, 54-79 mm FL). Coho salmon were also captured in the Taylor Creek drainage and were found up to Rm 880. Coho salmon made up 34% of the total catch in the Taylor Creek drainage ( $n = 72$ , mean FL 63.3 mm; range, 40-82 mm FL).

The only two juvenile Chinook salmon collected during our surveys were collected near the mouth of Taylor Creek. Both were collected in the lower 34 m of the stream.




FIGURE 2.7. —Distribution of juvenile coho salmon in Seattle’s streams, 2005-2006.

*Gasterosteidae, sticklebacks.*-- Threespine stickleback was one of the most wide-ranging species, occurring in one or more streams or ponds of each region except the Northwest region (Figure 2.8). They were present in seven drainages and were the only fish species present in Mapes Creek, Scheuerman Creek, and the upland ponds of South Fork Thornton Creek. In the Thornton Creek drainage, they were present from upland ponds of South Fork Thornton Creek to the mouth but were absent in North Fork Thornton Creek. At most stream sites, cutthroat trout were the dominant fish species present and threespine stickleback represented a small portion of the catch; however, at one 50 m reach in upper South Fork Thornton Creek (Rm 3,800) where only one cutthroat trout was collected, threespine stickleback were abundant ( $n = 78$ ; mean FL, 47.9 mm; range, 29-76 mm FL). In Scheuerman Creek, they were found in three ponds but were never collected in the stream. The lower 50 m of Taylor Creek was sampled in late June 2005 and 17 adult threespine stickleback (mean FL, 76.8 mm; range, 60-84 mm FL) were collected. Based on the sample date and their size, these were probably spawning adults from Lake Washington. The convergence pool of Thornton Creek with Lake Washington is probably also used as a spawning area for lake threespine stickleback, but because we sampled this area in late August we probably missed their spawning period.

Threespine stickleback appeared to be especially abundant in ponds and were usually associated with submerged aquatic vegetation. In streams, they were usually observed in quiet areas of pools. Proximity to a pond or lake appears to influence their abundance. For example, their abundance in the Thornton Creek system was much higher near Lake Washington and the upland ponds of the South Branch (Figure 2.8).

The size of threespine stickleback collected in ponds (mean FL, 35.3 mm) was generally smaller than the size collected in streams (mean FL, 43.7 mm)( $t$ -test,  $P < 0.001$ ). Generally, small fish  $< 30$  mm FL were not collected in streams. However, this may be partly because of differences in collection techniques. In streams, we primarily used electrofishing equipment and small threespine stickleback may be difficult to stun or may have been overlooked because of their small size. In ponds, we primarily used beach seines which may have adequately sampled all sizes of threespine stickleback. Alternatively, threespine stickleback may spawn primarily in the ponds and those observed in streams are subadults that have moved downstream from ponds. Also, streams often had

large predators such as cutthroat trout which may have consumed small threespine stickleback.




FIGURE 2.8. —Distribution of threespine stickleback in Seattle's streams and ponds, 2005-2006.

Cottidae, sculpins.-- In Thornton Creek, cottids were found as far upstream as the Meadowbrook Pond intake structure at Rm 2,000 (Figure 2.9). However, the weirs (Rm 400; 0.3 m vertical drop; Figure 2.10) at Sand Point Way appeared to act as a partial barrier to cottids. No prickly sculpin were found above the weirs but were found immediately below the weir. Coastrange sculpin were found above Sand Point Way but were significantly larger than those below the weir ( $t$ -test,  $P < 0.001$ ) and catch rates were substantially lower. Of the three sites above Sand Point Way where cottids were found, we only collected a total of nine coastrange sculpin. All coastrange sculpin above the weir were at least 87 mm TL (mean TL, 96.4 mm), while those below Sand Point Way ranged from 30 to 109 mm TL ( $n = 378$ ; mean TL, 52.2 mm). Thus, the weir appeared to be passable by only large coastrange sculpin. The ratio of prickly sculpin to coastrange sculpin tended to decrease at more upstream sites. Near the mouth in the convergence pool all cottids were prickly sculpin, but at 250-300 m from the mouth 38% were prickly sculpin, and at 350-400 m from the mouth 28% were prickly sculpin. These ratios were also influenced by the amount of pool/glide/riffle habitat available. In pools, 64% of the cottids were prickly sculpin, while they represented only 18% and 12% of the cottids in glides and riffles, respectively. Within the Thornton Creek basin, both prickly sculpin and coastrange sculpin were also found in Matthews Creek; prickly sculpin were in the small pond near the mouth and coastrange sculpin were in the creek above the pond.

Similar to Thornton Creek, the only cottid species found above a weir at Rm 42 (Figure 2.10; vertical drop, 0.3 m) on Piper's Creek was coastrange sculpin (mean TL, 64.4 mm; range, 38-97 mm TL) and they were significantly larger than coastrange sculpin below the weir (mean TL, 42.7 mm; range, 27-52 mm TL)( $t$ -test,  $P < 0.001$ ). Prickly sculpin and Pacific staghorn sculpin were found immediately below the weir but not above. At Taylor Creek, the upstream distribution of cottids appeared to be stopped by a short waterfall (0.3 m vertical drop; Figure 2.11) 63 m upstream from Lake Washington. Cottids were not collected immediately above this location or in several upstream locations.






FIGURE 2.9. —Distribution of cottids in Seattle’s streams and ponds, 2005-2006.

Of the cottids collected, 93% were coastrange sculpin ( $n = 27$ ; mean TL, 41.8 mm; range, 26-101 mm TL). The only cottid species in Washington Park Creek was prickly sculpin ( $n = 5$ ), which were found either at the mouth of the creek or 105 m upstream in a plunge pool at the base of a culvert (0.8 m vertical drop; Figure 2.12). The only cottid species observed in Fauntleroy Creek was Pacific staghorn sculpin, which were only observed near the mouth in the intertidal area. Cottids collected in Longfellow Creek were mostly large prickly sculpin ( $n = 9$ ; mean TL, 128 mm; range, 97-154 mm TL) and were only observed up to Rm 130. One juvenile Pacific staghorn sculpin was also collected.



FIGURE 2.10.-- Weirs in Thornton Creek (top panel) and Piper's Creek (lower panel) that appear to act as a partial barrier to upstream movement of cottids.



FIGURE 2.11.-- Short waterfall in lower Taylor Creek that appears to act as a complete barrier to upstream movement of cottids and juvenile salmonids. Several cottids were collected in the pool at the bottom of the photo but no cottids were ever collected upstream.



FIGURE 2.12.-- Culvert on Washington Park Creek that appears to act as a complete barrier to upstream movements of fish. Prickly sculpin and cutthroat trout were collected in the plunge pool shown but not upstream of the culvert.

Centrarchidae, sunfishes and black bass.-- Centrarchids observed in Seattle streams included smallmouth bass, largemouth bass, rock bass, and pumpkinseed. The only smallmouth bass (39 mm FL) collected was captured at the mouth of the Washington Park Creek. We assume that smallmouth bass only rarely use Seattle streams that drain into Lake Washington or the Lake Washington Ship Canal (LWSC). Rock bass, pumpkinseed, and largemouth bass were only found in the Thornton Creek watershed. In contrast to threespine stickleback, centrarchids were abundant in North Fork Thornton but absent in the South Fork Thornton Creek system, except for a few pumpkinseed in Kramer Creek. Adult rock bass and pumpkinseed and juvenile largemouth bass were primarily collected in a pond at the Jackson Park Golf Course. Pumpkinseed was the most numerous nonnative species captured. Additionally, we often collected several unidentified juvenile sunfish in Thornton Creek (including North Fork Thornton Creek), which we assumed were juvenile pumpkinseed.

Other introduced fish (Cyprinidae, Cobitidae, Ictaluridae).-- Introduced cyprinids observed in the study area included only goldfish and koi (an ornamental variety of common carp). Goldfish were found in three ponds, two in Washington Park and one in Discovery Park. Because a wide range of sizes were observed (range, 18-170 mm FL) in the Washington Park ponds, the goldfish populations appear to be self-sustaining through natural reproduction. Only one goldfish was collected in the Discovery Park and thus it's unclear if this represents a reproducing population. Koi were only observed in the Japanese Garden Pond in Washington Park and because they appear to be abundant and were present in a wide range of sizes, they probably are a naturally-producing population.

Two oriental weatherfish (family Cobitidae) were collected in the lower reach of Thornton Creek. Previous sampling has indicated they occur primarily in the LWSC, Lake Washington, and the Sammamish River (Tabor et al. 2001; E. Jeanes, R2 Resources, pers. comm.). Earlier sampling of Thornton Creek has also only found them in the lower reach (R. Tabor, unpublished data). Perhaps the weirs at Sand Point Way serve as a barrier to the upstream movement of this species.

Brown bullhead (family Ictaluridae) were collected at one location, a small pond in Washington Park. All fish appeared to be juveniles (range, 59-81 mm FL). Because we only made one seine haul in shallow water, we probably missed the adult fish.

### *Stream typing*

Results of stream typing are displayed in a series of maps in Appendix 1. Stream typing on the unnamed stream system #'s 1-4 were categorized as type 4 due to lack of fish presence and habitat characteristics (Appendix 1.1). The mainstem of Piper's Creek (stream system #5) from the mouth to the "twin pipes" (large culverts) at Rm 1,750 was classified as type 2 because of the stream size and high abundance of cutthroat trout (Appendices 1.2 and 1.3). Upstream to the headwaters of Piper's Creek was typed as type 3 along with most of Venema Creek, the lower section of Mohlendorph Creek, and the lower sections of two unnamed tributaries (Appendices 1.2 and 1.3). All other sections were classified as type 4. We did not collect any fish in Stream #8 but it was categorized as type 3 only because it met the habitat requirements for stream typing (Appendix 1.4). Streams system #'s 9-11 (Appendix 1.5) were typed as 5 whereas stream system #12(a-g; Appendix 1.6)) was classified as type 4. Licton Springs Park Creek was classified as type 3 based on habitat characteristics (Appendix 1.7).

Because of the stream size and high abundance of cutthroat trout, the lower mainstem of Thornton Creek and most of the North and South Branches were classified as type 2 (Appendices 1.8-1.16). The lower reaches of many tributaries were stream type 3. Upper reaches of these tributaries were stream type 4. Although no fish were found above the confluence of Maple Creek, or above 100<sup>th</sup> street on Willow Creek, or in Little's Creek, these streams were classified as type 3 base on habitat characteristics. Despite lack of fish, Inverness Creek (stream system #15) was also classified as type 3 due to habitat characteristics (Appendix 1.18). Ravenna Park Creek was categorized as type 3 (Appendix 1.18).

With the west-central region, Wolfe Creek, Scheuerman Creek, and a small unnamed tributary to Scheuerman Creek were typed as 3, while all other streams in this region were type 4 or 5 (Appendices 1.19 and 1.20).

Washington Park Creek was the only creek that was a type 3 in the east-central region, while all other creeks were type 4 or 5 (Appendices 1.21 and 1.22).

The mainstem of Longfellow Creek was typed as type 3 with three small tributaries typed as 4 (Appendices 1.23 and 1.24). Puget Creek was a type 4 stream except for a small section that was type 3 (Appendix 1.25). Although we did not find any fish in Schmitz Creek, its mainstem was still a type 3, with several small tributaries as type 4, due to adequate habitat characteristics (Appendix 1.26). Stream #'s 39 (Mee Kwa Mooks Park), 40, and 41 were all typed as 4 (Appendix 1.27). Fauntleroy Creek mainstem was all type 3 (Appendix 1.27), and Seola Beach Creek was a type 5 (Appendix 1.28). The different sections of the Durham Creek system were typed as 3 and 4, but some sections could not be surveyed due to inaccessibility (Appendix 1.28).

Of the southeast region streams, Mapes Creek and Taylor Creek were typed as 3 while all other creeks were classified as type 4 (Appendices 1.29 and 1.30).

## Discussion

Several streams and ponds were void of fish. Generally, the lack of fish appeared to be due to one or more of three possible explanations: 1) small headwater streams with little available habitat, 2) presence of barriers and loss of connectivity to fish-bearing systems, and 3) severely degraded habitat conditions. Many of the small streams we examined that were void of fish had streamflows less than 0.03 cfs and the average wetted width was less than 0.7 m. Even under ideal conditions, streams of this size would probably be void of fish most of the year. Latterell et al. (2003) found the upstream extent of trout in western Cascade Mountains was influenced by gradient, stream size, and pool availability. In most Seattle's streams, trout distribution did not appear to be constrained by gradient; however, stream size and pool availability may be important factors. In a study of 79 headwater streams in southwestern Washington, Fransen et al. (1998) found streams needed to have a mean annual flow of at least 0.5 cfs to have the potential to produce the minimum amount of food needed to sustain a fish. Although we only sampled in the summer low-flow period, many of small streams we surveyed were unlikely to meet this criteria. Alternatively, cutthroat trout and juvenile coho salmon may inhabit streams as small as 1.2 m bankfull width (Rosenfeld et al. 2000) and thus many of the Seattle's fishless streams may be large enough to support fish. Also, fish-bearing streams in the western Cascade Mountain area are generally present in watersheds that are at least 22 ha (Trotter 2000). On a cursory examination of watershed sizes, many of the fishless streams in Seattle area appear to occur in watersheds that are large enough to support fish-bearing streams. Perhaps under pristine conditions, many of these fishless streams had fish populations.

At some locations, such as Willow Creek, Maple Creek, and Washington Park Creek, a steep or perched culvert appeared to prevent fish from utilizing available upstream habitat. Additionally, in some systems (i.e., Durham Creek, Schmitz Creek, and Puget Creek) the lower section of the stream is in a long culvert and its unclear if the culvert is a barrier to fish movements. Fish populations above the culverts

The habitat conditions of urban streams including Seattle's streams are often severely degraded. The number, size, and depth of pools are greatly reduced. For example, Schmitz Creek had reasonable streamflow but few pools were present and those that were present had maximum depths less than 0.2 m, which provide little cover for juvenile salmonids.

In general, our results were consistent with Wild Fish Conservancy results (Washington Trout 2000). The distribution of most fish including cutthroat trout and sculpins appears to be quite similar between the two studies. However, there were a few notable differences. One major difference was the difference of fish distribution in Venema Creek. Wild Fish Conservancy observed fish up to the headwaters (approximately Rm 900), while we only observed them up to a logjam at Rm 320. A small landslide (approximately Rm 700) had occurred in this drainage a few months before our survey, which may have extirpated trout from the upper reaches. Another major difference was the upstream extent of fish in Taylor Creek. We observed fish up to Rm 1,100, while Wild Fish Conservancy (Washington Trout 2000) only observed fish up to a perched culvert at Rm 450. Apparently, the culvert, which was replaced in 1999, allowed fish to move upstream. Additionally, we conducted winter surveys and were able to document fish use in Puget Creek as well as upstream of the twin culverts on Piper's Creek, and upper reach of South Branch Thornton Creek. Wild Fish Conservancy (Washington Trout 2000) observed juvenile Chinook salmon in Piper's Creek and Thornton Creek, while we only observed one Chinook salmon at the mouth of Taylor Creek. Wild Fish Conservancy surveys were conducted from late June to mid July, 1999. We may have missed the occurrence of juvenile Chinook salmon in many streams because many of our surveys were done after July when the vast majority of juvenile Chinook salmon had already outmigrated.

Previous studies have found that as watersheds become more urbanized the ratio of juvenile coho salmon abundance to cutthroat trout abundance decreases (Scott et al. 1986; Serl 1999; Horner and May 1998; Seiler et al. 2005). Horner and May (1998) found the coho/cutthroat ratio is greater than 2:1 in watersheds with low levels of development and



0.5/1 in highly urbanized streams. Thornton Creek and Piper's Creek systems appear to fit this pattern for urbanized stream systems. The ratios were 0.016:1 for Thornton Creek and 0.048:1 for Piper's Creek. Taylor Creek had a much higher ratio (1.24:1) but it is unclear why. The effects of urbanization may be less severe on this system. Also, Taylor Creek is a small system and the number of spawning adults may vary greatly from year to year. We are uncertain about the origin of coho salmon which may have been naturally produced or stocked hatchery fish.

In contrast to Piper's Creek, Thornton Creek, and Taylor Creek, cutthroat trout appear to be extremely rare (one cutthroat trout collected from approximately 500 m of stream length sampled) in Longfellow Creek despite a large amount of available habitat (Figure 2.13). Typically, cutthroat trout occupy a wide-range of habitats in western Washington. They are usually found further upstream than other fish species (Trotter 2000) and are abundant in Puget Sound and lowland lakes like Lake Washington. One possibility for the lack of cutthroat trout in Longfellow Creek is the long culvert at the lowest stream section which may serve as a barrier. However, it would seem unlikely that the culvert is a hydrologic or visual barrier to cutthroat trout since adult coho salmon as well as cottids can move upstream through the culvert and there are a number of skylights along its 700-m course. The stream appears to be more turbid than other streams but it's doubtful if this would dramatically reduce their abundance. The lack of cutthroat trout may be related to contaminants. High rates of pre-spawning mortality of adult coho salmon have been observed in Longfellow Creek and researchers believe it's likely caused by exposure to environmental contaminants (Reed et al. 2004). Whether contaminants may affect cutthroat trout rearing, spawning, or incubating in Longfellow Creek is unknown.



FIGURE 2.12.—Photo of Longfellow Creek at rm 2,900, August 2, 2005. In this 50 m reach, only three juvenile coho salmon and two rainbow trout were collected. Streams with similar habitat conditions in other regions of Seattle would be expected to have large numbers of cutthroat trout present.

The lack of cutthroat trout was also evident in other southwest streams. Juvenile coho salmon or rainbow trout were collected in Fauntleroy Creek, Puget Creek, and Durham Creek but cutthroat trout were never collected. Similar to Longfellow Creek, each of these streams does have a culvert close to its estuary but it's unclear if they are barriers

to cutthroat trout. The lack of cutthroat trout may be related to the size of the creek. However, Venema Creek and Kramer Creek are similar-sized creeks and cutthroat trout were abundant. The level of environmental contaminants in these creeks is not known. Additional research is needed to better understand why cutthroat trout are rare in the southwest streams including Longfellow Creek but abundant in other similar-sized streams.

At some locations, we were unclear if the observed fish were part of a self-sustaining population or were simply recently stocked fish. This was particularly true for juvenile coho salmon. At Fauntleroy Creek, Durham Creek, and Piper's Creek, members of local enhancement groups indicated coho salmon were often stocked. A few rainbow trout were collected in Ravenna Creek and its unclear how these fish got there. Most likely they were stocked fish. A single goldfish was collected in a Discovery Park pond and may have been recently stocked by a local resident.

Besides the Washington Trout (2000) surveys, there has been few comprehensive surveys of Seattle's stream except in Thornton Creek and Piper's Creek. Results of surveys from Thornton Creek (Muto and Shefler 1983; Ludwa et al. 1997; Leavy et al. 2007) found the same general fish distribution as we observed. In all studies, cutthroat trout was the dominant species collected, the cutthroat trout to juvenile coho salmon ratio was low, and sculpin were only collected in sites close the creek mouth.

Comprehensive surveys of Piper's Creek were conducted by Pfeifer (1984) in 1984 and by Thomas (1992) in 1991 and 1992. Results of those studies and this study are summarized in Table 6.1. There were three notable differences between the studies. First, the number of cutthroat trout/stream length we collected was substantially higher than the other two studies. We collected our samples in October while the other studies were conducted in July or May; however we would expect the trout population to be lower in October as the number of age-0 trout is reduced over the over the course of the summer. Secondly, the distribution and abundance of cottids appears to have been reduced from 1984 to 2005. Thirdly, Thomas (1992) was the only study to document the presence of Pacific giant salamanders *Dicamptodon tenebrosus*.

The increase of cutthroat trout abundance may be due to several factors including stream restoration efforts, reduction in angling effort, and reduction in the abundance of other fish and salamanders. In the summer of 1991, log and rock weirs were installed, which increased the pool-to-riffle ratio from 0.13 to 0.29 (Thomas 1992). Additionally, several K-weirs were installed in 1998 and modified in 1999. The creation of larger, deeper, and more frequent pools may have greatly increased the abundance of cutthroat trout. Also, competition and predation by sculpin and Pacific giant salamanders could have reduced the abundance of cutthroat trout in the 1980's and 1990's. Additionally, hatchery rainbow trout were sometimes stocked into upper Piper's Creek in the 1980's, which may have competed or possibly displaced some cutthroat trout. In 2005, the abundance of competitors and predators of cutthroat trout appears to be reduced.

Pfeifer (1984) found cottids were abundant in both areas he sampled in the lower reach of Piper's Creek. In contrast, Thomas (1992) and this study found few cottids were present in the lower reach. Currently, a small weir is located just upstream of the railroad culvert that may limit recruitment of young cottids to the lower reach. The date this weir was constructed is unknown.

Combined, Thomas (1992) collected 16 Pacific giant salamanders in 1991 and 1992. Pfeifer (1984) did not document the presence of Pacific giant salamanders but he may have missed them because he only sampled the lower 1.0 km and Thomas (1992) found they were primarily upstream of this location. We conducted several surveys throughout the basin and salamanders were never observed. Pacific giant salamanders have been shown to be sensitive to increased sedimentation (Welsh and Ollivier 1998). Also, substrate conditions (high embeddedness and few large substrates) and the lack of large woody debris in Piper's Creek may have affected salamander abundance. The cumulative effects of urbanization may have caused salamanders to be extirpated from this system.

TABLE 2.10. -- Electrofishing results of three studies (Pfeifer 1984, Thomas 1992, and this study) of Piper's Creek. The month and year are when the stream was sampled. The lower reach extends from the railroad culvert (Rm 0.0) to the sewage treatment culvert (Rm 650) and the upper reach is from the sewage treatment culvert to the twin pipes (Rm 1,750). Pfeifer (1984) only sampled up to the sewage treatment culvert, while Thomas (1992) sampled up to the twin pipes. For this study, we only used data from sites that would match the other studies. Pfeifer (1984) used multiple-pass electrofishing whereas single-pass electrofishing was used by the other studies. We used data from Pfeifer's first pass to make comparisons to the other studies.

| Study   | Month, Year | Reach | Distance surveyed (m) | Catch     |      |         | Number per stream length |      |         | Percent of catch |       |         |
|---------|-------------|-------|-----------------------|-----------|------|---------|--------------------------|------|---------|------------------|-------|---------|
|         |             |       |                       | Cutthroat | Coho | Cottids | Cutthroat                | Coho | Cottids | Cutthroat        | Coho  | Cottids |
| Pfeifer | July-84     | Lower | 66                    | 8         | 15   | 57      | 0.12                     | 0.23 | 0.86    | 10.00            | 18.75 | 71.25   |
| Thomas  | May-91      | Lower | 600.6                 | 56        | 32   | 39      | 0.09                     | 0.05 | 0.06    | 44.09            | 25.20 | 30.71   |
| Thomas  | May-92      | Lower | 600.6                 | 395       | 16   | 86      | 0.66                     | 0.03 | 0.14    | 79.48            | 3.22  | 17.30   |
| Lantz   | October-05  | Lower | 132                   | 132       | 10   | 13      | 1.00                     | 0.08 | 0.10    | 85.16            | 6.45  | 8.39    |
| Thomas  | May-91      | Upper | 419.5                 | 80        | 0    | 0       | 0.19                     | 0.00 | 0.00    | 100              | 0     | 0       |
| Thomas  | May-92      | Upper | 419.5                 | 157       | 0    | 0       | 0.37                     | 0.00 | 0.00    | 100              | 0     | 0       |
| Lantz   | October-05  | Upper | 100                   | 90        | 0    | 0       | 0.90                     | 0.00 | 0.00    | 100              | 0     | 0       |

Although cottids are an abundant, widespread group of fish in the Pacific Northwest, they have a limited distribution in Seattle streams. They were present in six drainages and were usually only found in the lower reaches, often only a few meters from Lake Washington or Puget Sound. The cottids found within the streams of WRIA 8 (Lake Washington basin and a few nearby small independent drainages) can be divided within three types: 1) estuarine species, 2) lowland freshwater species, and 3) upland freshwater species. Although there is often a large degree of overlap, these groups generally occupy different areas of a basin. Estuarine species are primarily found in the lower sections of streams that are under tidal influence; lowland freshwater species are widespread in lowland lakes and usually found in the lower reaches of streams and rivers; and upland freshwater species are found in the middle and upper reaches of streams and rivers and upland lakes such as Chester Morse Lake.

The vast majority of estuarine cottids were Pacific staghorn sculpin. Young Pacific staghorn sculpin often move upstream in the spring and can tolerate areas with low salinity (Wydoski and Whitney 2003). The farthest upstream we collected a Pacific staghorn sculpin (44 mm TL) was in Longfellow Creek at river kilometer 0.8. Other estuarine

cottids, such as sharpnose sculpin, occasionally enter freshwater but their numbers are usually quite low in comparison to Pacific staghorn sculpin.

Lowland freshwater cottids in the Lake Washington basin consist of coastrange sculpin and prickly sculpin. Both species have planktonic larvae, relatively small eggs, and have higher fecundity rates than most other freshwater cottids (Wydoski and Whitney 2003). They migrate downstream in the spring to spawn, and after spawning, adults and juveniles migrate upstream in the summer and fall (Morrow 1980). Additionally, both species inhabit lacustrine and estuarine environments.

Because the lowland sculpin species typically spawn in lower stream reaches and are not strong swimmers, their ability to disperse to upstream habitats can be limited by small barriers. For some sculpin species, such as the European bullhead *C. gobio*, impassable barriers may be as low as 18-20 cm (Utzinger et al. 1998). Mason and Machodori (1976) found obstructions 30 cm high were impassable to prickly sculpin and 45 cm high were impassable to coastrange sculpin. Shapovalov and Taft (1954) suggested that low-head dams (approximately 90 cm high) were an effective method of eliminating upstream populations of prickly sculpin and coastrange sculpin. In a recent study of sculpin barriers in northwest Washington, LeMoine (2007) found that perched culverts and fish ladders often limited the upstream distribution of prickly sculpin and coastrange sculpin. Our results appear to be consistent with these studies. The upstream distribution of prickly sculpin appeared to be stopped by barriers on Thornton Creek, Taylor Creek, Piper's Creek, and Washington Park Creek. Likewise, the upstream distribution of coastrange sculpin was stopped by the same barrier on Taylor Creek; however, a few coastrange sculpin were found above barriers on Thornton Creek and Piper's Creek. These fish were mostly large individuals, suggesting the barriers limited the upstream movement of small coastrange sculpin but some large individuals were able to move upstream of the barriers probably because of their superior swimming ability.

Upland freshwater cottids in the Lake Washington basin consist of riffle sculpin *C. gulosus*, torrent sculpin *C. rhotheus*, and shorthead sculpin *C. confusus* (Tabor et al.

2007a). These species have larger eggs and lower fecundity than the lowland species (Wydoski and Whitney 2003). Larvae can be either demersal (riffle sculpin) or planktonic (shorthead sculpin). None of the upland species were observed in Seattle streams. Shorthead sculpin typically inhabit coldwater streams and thus would not be expected to inhabit Seattle's streams. However, riffle sculpin and torrent sculpin are widespread in tributaries to Lake Washington and Lake Sammamish and thus would be expected to occur in some Seattle streams such as Thornton Creek. For example, riffle sculpin and torrent sculpin are common in May Creek, a similar-sized stream as Thornton Creek. Reasons why the upper reaches of Thornton Creek and other streams are void of upland cottids is unclear. Increased urbanization often leads to severe changes in stream habitat conditions such as higher peak flows, elevated water temperatures, reduction in water quality and reduction of habitat quality (i.e., reduction in the amount of woody debris and pools). The effects of high peak flows may be especially deleterious to cottid populations (Erman et al. 1988).

If cottid populations have been extirpated and habitat conditions improve, coastrange sculpin and prickly sculpin, which are widespread in Lake Washington and in nearshore areas of Puget Sound, can easily move into the lower stream reaches. However, their upstream distribution will be limited by barriers. Upland cottid species may have a difficult time returning to many streams once they have been extirpated. They often have a restricted home range and disperse slowly. For example, Moyle (2002) noted that riffle sculpin in a small stream California took over 18 months to recolonize a riffle that went dry that was 500 m downstream of a large population. The upland cottid species have not been documented in Puget Sound or Lake Washington. Large lakes such as Lake Washington may also serve as a barrier to dispersal to other streams due to predation from prickly sculpin and exotic fishes such as yellow perch *Perca flavescens* and smallmouth bass.

Attempts to rehabilitate urban streams should consider the reintroduction of upland cottids. In particular, riffle sculpin and torrent sculpin would be good candidates for a recolonization experiment. These species can complete their life-cycle in a relatively small area. Reintroduction of sculpin would provide valuable information on stream habitat

conditions and help identify factors affecting their survival. These types of experiments may provide valuable information on why more urbanized systems tend to have few sculpin (Matzen and Berge 2008). Freshwater sculpin have been used as an indicator species of stream health (Gray et al. 2004; Gray et al. 2005; Adams and Schmetterling 2007). Groups of sculpin could be introduced into different streams and different habitat types to understand factors affecting their survival. Locations with both stream and pond habitat such as Littles Creek in Jackson Park Golf Course would have diverse habitat conditions where cottid populations would be more likely to withstand high-flow conditions. The introduced sculpin should be PIT tagged and their movements monitored with mobile and stationary receivers to understand factors (i.e. high flow events) related to their movements. Additionally, sculpin in other less-developed watersheds such as May Creek could be PIT tagged and their movements compared to sculpin movements in Thornton Creek.

Introduction of any species should be considered carefully before any decision is made. Impacts to the existing ecosystem can often be difficult to predict, particularly in systems that have a complex food web. Impacts of introduced sculpin to the ecosystem would likely be due to predation and/or competition with native salmonids. Sculpin have been documented to prey on juvenile salmonids and salmonid eggs in many systems (Foote and Brown 1998; Tabor et al. 2007b); however, sculpin do not appear to be a major predator in most cases (Moyle 1977). Sculpin may also compete with salmonids for benthic macroinvertebrates. The degree of competition can vary between size and species of salmonid present (Zimmerman and Vondracek 2007). Alternatively, sculpin may provide a valuable prey resource for stream-dwelling salmonids. Consumption of small sculpin by cutthroat trout and other salmonids has been documented in Pacific Northwest streams and rivers (Lowry 1966; Price 2006). Exactly how sculpin would affect the food web and salmonid populations is difficult to predict. If sculpin are introduced, monitoring of salmonid and sculpin diet and growth and the macroinvertebrate community will be needed to understand their interrelationships.



Managers should also consider removing man-made barriers that limit the upstream movement of coastrange sculpin and prickly sculpin. Currently, barriers on Piper's Creek, Thornton Creek, and Washington Park Creek appear to restrict upstream movement of these cottids. Monitoring efforts before and after the barriers are removed would provide information on movements of prickly sculpin and coastrange sculpin into upstream reaches.

In addition to constraints to upstream movements of cottids, small barriers may also restrict the distribution of juvenile salmonids. Juvenile Chinook salmon and coho salmon often move into non-natal tributaries to use as rearing areas, sometimes for several months. Juvenile cutthroat trout may also move into small streams but it is difficult to determine whether a stream is a natal or non-natal stream because adults can spawn in small streams. Because the abundance of juvenile coho salmon and Chinook salmon is low in Seattle's streams it is usually difficult to assess whether a particular obstacle is a barrier. Little information is available on what constitutes a barrier to juvenile salmonids. In earlier surveys of Matthews Creek (Tabor et al. 2004), we observed several juvenile coho salmon at the base of a log weir and they did not appear to move upstream until the lake level rose and the weir was no longer a barrier. Juvenile Chinook salmon, however, were able to move upstream past the barrier even when the lake level was low. In Taylor Creek, juvenile Chinook salmon have only been observed up to the small waterfall at Rm 42. Presumably this waterfall, that is a barrier for cottids, is also barrier for juvenile Chinook salmon. However, the stream section between

Across their native distribution, three-spine sticklebacks inhabit a vast array of habitat types, from small streams to large lakes to the marine environment including the open ocean (McPhail 2007). Both resident and anadromous forms occur. Typically, they are found in the lower reaches of coastal streams and occur upstream to major fish barriers such as waterfalls. In some areas, they have been introduced, probably as a result of contamination with some type of fish transfer (i.e. trout plantings or baitfish). In Seattle's lakes, ponds, and streams, they also inhabit a large variety of habitat types. They are an important component of the pelagic zone in Lake Washington. They are widespread throughout the South Branch of Thornton Creek. In the Scheuerman Creek

system and Mapes Creek, they were the only species present. They were also abundant in several ponds and appeared to be closely associated with aquatic macrophytes. They are native to Lake Washington but it is unclear if their distribution in Seattle's streams and ponds is a part of the native distribution or is result of introductions.

Nonnative fish species were collected in some streams; however their abundance was generally low in comparison to native fish species. The species of nonnative fish in Seattle's streams are not well-adapted to inhabit small streams and do not appear to be a significant threat to the ecosystem health. Most nonnative fish collected were juvenile sunfish from upstream ponds. In fact they may be somewhat beneficial; as juvenile sunfish move out of pond habitats they may become a food source for native salmonids such as cutthroat trout. Another species, oriental weatherfish was collected in lower Thornton Creek as early as 1998 (Resource Planning Associates 1998), yet there does not appear to be any large increase in their abundance.

In some ponds, nonnative species including largemouth bass, rock bass, pumpkinseed, and brown bullhead may dominate the fish fauna. In these habitats they could impact the abundance of native fishes through predation and perhaps competition. If native fishes inhabit the pond throughout the year (threespine stickleback and sculpin) or are only present when water temperatures are cool (cutthroat trout and juvenile coho salmon), they could have some overlap with nonnative fishes and could be negatively impacted. Additionally, amphibians, crayfish, and other pond organisms may be impacted by introductions of nonnative fishes.

## CHAPTER 3. REFERENCE SITE FISH DENSITIES

### Introduction and Methods

A key element of management of Seattle's streams is long-term monitoring, which includes effectiveness monitoring of stream restoration projects and long-term monitoring to assess changes in ecosystem health. Within each of the five major watersheds, we undertook more intensive sampling of the fish community at one or more sites. These sites will serve as reference sites, which will be sampled again and again to assess long-term changes in the fish community. Sites in Thornton Creek watershed were selected because they are restoration sites (Table 3.1). Instream restoration work of the Maple Leaf Reach of the South Branch Thornton Creek was completed in 2008, which included the addition of two logjams and some large boulders. We conducted pre-project monitoring of this site in 2005 and 2006. Within the Thornton Creek basin, we also conducted pre-project monitoring of the lower section of Kramer Creek. The city plans to reconfigure the convergence area and stream channel; however, the exact project plan and schedule is not known at this time. Sites in Piper's Creek will be used to assess upland land-use changes to reduce stormwater flows and create a more natural drainage pattern. Sites in the other three major watersheds will serve as long-term reference sites to assess ecosystem health. Our reference site in Taylor Creek is above a man-made barrier, that the city plans to be removed, but it may be several years before the barrier is removed. Our sampling will provide baseline information once the barrier is removed. One site was surveyed on each stream, except Venema Creek where two sites were surveyed, one below the confluence with Mohlendorph Creek and one above the confluence. Reference sites were between 50 and 130 m long.

We used depletion procedures to determine the overall fish abundance and biomass in each site. Multiple-pass electrofishing was conducted to capture fish. We first divided each site into habitat units: riffles, pools, and glides. Block nets (5.8-m long by 1.2-m high with 5-mm square mesh) were used to isolate each habitat unit. Once all the nets were in place, crews waited approximately 15 minutes before sampling. Depending on the size of the reach, two or three crew members worked together to

TABLE 3.1. Location (Rm, river kilometer) and sample dates of reference sites surveyed within Seattle's streams. Appendix number is the map which displays the location of each sample site (sample #).

| Stream system # | Site location               | Rm      | Date Sampled | Sample # | Appendix |
|-----------------|-----------------------------|---------|--------------|----------|----------|
| 5               | Piper's Creek               | 820-870 | 3-Mar-05     | 500      | 1.2      |
|                 |                             |         | 19-Oct-05    | 501      | 1.2      |
|                 |                             |         | 29-Sep-06    | 502      | 1.2      |
|                 | Venema Creek - lower        | 0-50    | 3-Mar-05     | 510      | 1.2      |
|                 |                             |         | 19-Oct-05    | 511      | 1.2      |
|                 |                             |         | 29-Sep-06    | 512      | 1.2      |
|                 | Venema Creek - upper        | 200-250 | 3-Mar-05     | 513      | 1.2      |
|                 |                             | 270-330 | 20-Oct-05    | 514      | 1.2      |
|                 |                             |         | 29-Sep-06    | 515      | 1.2      |
|                 | Mohlendorph Creek           | 100-150 | 19-Oct-05    | 516      | 1.2      |
| 14              | South Branch Thornton Creek | 550-670 | 24-Feb-05    | 520      | 1.10     |
|                 |                             |         | 13-Oct-05    | 521      | 1.10     |
|                 |                             |         | 2-Aug-06     | 522      | 1.10     |
|                 | Kramer Creek                | 0-110   | 25-Feb-05    | 523      | 1.10     |
|                 |                             |         | 28-Sep-05    | 524      | 1.10     |
|                 |                             |         | 26-Jul-06    | 525      | 1.10     |
| 34              | Longfellow Creek            | 100-170 | 7-Aug-06     | 530      | 1.23     |
| 42              | Fautleroy Creek             | 220-320 | 18-Oct-05    | 540      | 1.27     |
|                 |                             |         | 8-Aug-06     | 541      | 1.27     |
| 49              | Taylor Creek                | 430-530 | 17-Oct-05    | 550      | 1.30     |
|                 |                             |         | 3-Aug-06     | 551      | 1.30     |

collect fish from each habitat. One person was used to operate the electrofisher and one or two were needed to net the stunned fish. Electrofishing was conducted with pulsed DC current set at 200 to 300 volts depending on the stream conductivity and effectiveness of stunning the fish. Sampling began at the downstream end. At least three passes were conducted. We stopped sampling on the third pass if the catch (number of fish) on the third pass was less than 50% of the catch on the second pass. If a 50% depletion of fish was not achieved on the third pass for any habitat, a fourth or sometimes a fifth pass was conducted until a 50% depletion was achieved. Fish from each pass and each habitat type were kept in separate buckets. Fish were anesthetized with MS-222 and identified. The fish length (nearest mm) and weight (nearest 0.1 g) were measured. After fish had been processed and released and the block nets removed, we conducted a habitat survey of the reference site. Habitat data collected included:

Habitat length- Stream length from the downstream end of the habitat to the upstream end via the thalweg of the stream.

Habitat width- The wetted width of the habitat was calculated from the mean of three equidistant width measurements:  $\frac{1}{4}$ ,  $\frac{1}{2}$ , and  $\frac{3}{4}$  of the stream length of each habitat unit.

Depth- A mean and maximum depth was recorded at each habitat unit. The average depth was measured by randomly placing the hand held stadia rod at different locations within the habitat and averaging those values. The maximum depth of the habitat was collected by using the stadia rod to find the deepest part of each habitat unit. For pools we also measured the depth of the tailout or pool outlet. The tailout measurement is taken at the shallowest part (lengthwise) of the pool and the deepest part of the tailout (cross-section). The residual pool depth is the maximum pool depth minus the tailout depth.

Large woody debris- At each habitat unit, LWD was counted. A piece of large woody debris was any wood that was over 2 m in length and was 10-cm wide at the midpoint and was in contact with the stream. Procedures followed the TFW methodology for Level 1 (Zone 1) surveys of LWD (Schuett-Hames et al. 1999).

Estimated population ( $\hat{N}$ ) size for each reference site was calculated using the procedures of Carl and Strub (1978). A single population estimate was made for each site. We did not make a estimate for each habitat unit because the number of fish in each habitat unit was usually too low to make a precise population estimate.

## Results

*Piper's Creek watershed*.-- Piper's Creek reference site was sampled on three occasions: March 3, 2005, October 19, 2005, and September 29, 2006. At this location, the mean stream width was 2.8 m, maximum depth was 0.5 m, and the habitat consisted of approximately 26% pools (by area), 48% riffles and 26% glides (Table 3.2). Ninety-nine percent (191 of 193) of the fish captured at this site were cutthroat trout. The other two fish were coho salmon; one appeared to be a returning adult and was not used in our density estimates. The estimated population size ( $\hat{N}$ ) of cutthroat trout was similar between March 2005 ( $\hat{N} = 43$ ) and October 2005 ( $\hat{N} = 44$ ), but was substantially higher in September 2006 ( $\hat{N} = 117$ ; Table 3.3). In September 2006, 77% of the cutthroat trout were  $< 100$  mm FL; whereas in March 2005 and October 2005, 47% and 29%, respectively were  $< 100$  mm FL (Figure 3.1).

TABLE 3-2.-- Habitat information and number of fish collected in different habitat types of reference sites in the Piper's Creek watershed. %Fish is the percent of fish for each reference site that was collected in each habitat type (not adjusted by area). NA = no data available.

| Site location   | Date Sampled    | Habitat Types | Number of Habitats | Combined Lengths (m) | Mean Wetted Width (m) | Area (m <sup>2</sup> ) | Mean Depth (m) | Maximum Depth (m) | # of Fish Captured | % Fish |
|-----------------|-----------------|---------------|--------------------|----------------------|-----------------------|------------------------|----------------|-------------------|--------------------|--------|
| Mainstem        | 3-Mar-05        | Riffle        | 2                  | 35.5                 | 2.8                   | 99.4                   | 0.10           | 0.17              | 21                 | 56.8   |
|                 |                 | Pool          | 1                  | 2.5                  | 1.8                   | 4.5                    | 0.20           | 0.50              | 5                  | 13.5   |
|                 |                 | Glide         | 1                  | 10.0                 | 3.1                   | 31.0                   | 0.15           | 0.20              | 11                 | 29.7   |
|                 | 19-Oct-05       | Riffle        | 4                  | 24.5                 | 2.1                   | 51.5                   | 0.15           | 0.30              | 13                 | 30.2   |
|                 |                 | Pool          | 2                  | 17.5                 | 3.4                   | 59.5                   | 0.20           | 0.45              | 20                 | 46.5   |
|                 |                 | Glide         | 2                  | 9.5                  | 4.6                   | 43.7                   | 0.20           | 0.40              | 10                 | 23.3   |
|                 | 29-Sep-06       | Riffle        | 3                  | 19.5                 | 2.6                   | 50.8                   | 0.09           | 0.10              | 18                 | 15.9   |
|                 |                 | Pool          | 4                  | 19.2                 | 2.5                   | 48.0                   | 0.30           | 0.42              | 64                 | 56.6   |
|                 |                 | Glide         | 1                  | 11.1                 | 3.2                   | 35.5                   | 0.19           | 0.28              | 31                 | 27.4   |
| Lower Venema Cr | 3-Mar-05        | Riffle        | 5                  | 41.6                 | 1.4                   | 58.2                   | 0.10           | 0.15              | 17                 | 58.6   |
|                 |                 | Pool          | 4                  | 9.5                  | 2.1                   | 20.0                   | 0.20           | 0.47              | 12                 | 41.4   |
|                 | 19-Oct-05       | Riffle        | 6                  | 40.2                 | 1.2                   | 48.2                   | 0.05           | 0.15              | 3                  | 25.0   |
|                 |                 | Pool          | 4                  | 13.1                 | 1.9                   | 24.9                   | 0.25           | 0.38              | 7                  | 58.3   |
|                 |                 | Glide         | 1                  | 2.5                  | 1.3                   | 3.3                    | 0.10           | 0.61              | 2                  | 16.7   |
|                 | 28-Sep-06       | Riffle        | 8                  | 37.3                 | 1.3                   | 48.5                   | 0.08           | NA                | 12                 | 27.9   |
|                 |                 | Pool          | 7                  | 16.4                 | 1.8                   | 29.5                   | NA             | 1.80              | 31                 | 72.1   |
|                 | Upper Venema Cr | 3-Mar-05      | Riffle             | 5                    | 41                    | 1.5                    | 61.5           | 0.17              | 0.21               | 3      |
| Pool            |                 |               | 4                  | 8                    | 1.5                   | 12.0                   | 0.31           | 0.39              | 4                  | 57.1   |
| 20-Oct-05       |                 | Riffle        | 5                  | 19.9                 | 1.2                   | 23.9                   | 0.05           | 0.16              | 0                  | 0.0    |
|                 |                 | Pool          | 2                  | 3                    | 1.7                   | 5.1                    | 0.13           | 0.30              | 2                  | 28.6   |
|                 |                 | Glide         | 3                  | 9                    | 1.7                   | 15.3                   | 0.11           | 0.22              | 4                  | 57.1   |
| 29-Sep-06       |                 | Cascade       | 1                  | 18                   | 2.8                   | 50.4                   | 0.04           | 0.20              | 1                  | 14.3   |
|                 |                 | Riffle        | 4                  | 38.4                 | 1.2                   | 46.1                   | 0.05           | NA                | 3                  | 42.9   |
|                 |                 | Pool          | 2                  | 2.25                 | 1.5                   | 3.4                    | NA             | 0.25              | 1                  | 14.3   |
| 19-Oct-05       |                 | Cascade       | 1                  | 10.97                | 0.8                   | 8.8                    | 0.06           | NA                | 3                  | 42.9   |
|                 | Riffle          | 8             | 44.7               | 0.88                 | 15.9                  | 0.02                   | 0.11           | 0                 | --                 |        |
| Mohlendorph Cr  | 19-Oct-05       | Glide         | 8                  | 15.9                 | 1.08                  | 44.7                   | 0.06           | 0.20              | 0                  | --     |

TABLE 3.3.-- Population and density estimates of reference sites in Piper's Creek and Thornton Creek watersheds. Data is for cutthroat trout except data in parentheses which is for all fish combined. Other fish included rainbow trout, coho salmon, threespine stickleback, and pumpkinseed. Population estimates were calculated using depletion techniques of Carl and Strub (1978). Biomass (fish (g)/m<sup>2</sup>) is calculated by multiplying Fish/m<sup>2</sup> by the average weight (g) for each species. No fish were collected in Mohlendorph Creek and it is not included in this table.

| Watershed      | Site location     | Date sampled | Estimated population | Mean length (mm) | Mean weight (g) | Fish / m    | Fish / m <sup>2</sup> | Fish (g) / m <sup>2</sup> | Length of unit (m) | Mean wetted width (m) | Area (m <sup>2</sup> ) |
|----------------|-------------------|--------------|----------------------|------------------|-----------------|-------------|-----------------------|---------------------------|--------------------|-----------------------|------------------------|
| Piper's Creek  | Mainstem          | 3-Mar-05     | 43                   | 119.7            | 20.50           | 0.90        | 0.32                  | 6.56                      | 48.0               | 2.8                   | 134.9                  |
|                |                   | 19-Oct-05    | 43                   | 113.9            | 21.10           | 0.83        | 0.28                  | 5.91                      | 51.5               | 3.0                   | 154.5                  |
|                |                   | 29-Sep-06    | 116 (117)            | 91.7             | 11.60           | 2.35 (2.35) | 0.87 (0.87)           | 10.09 (10.15)             | 49.8               | 2.7                   | 134.5                  |
|                | Venema Cr (lower) | 3-Mar-05     | 29                   | 113.1            | 21.80           | 0.57        | 0.37                  | 8.07                      | 51.1               | 1.5                   | 78.2                   |
|                |                   | 19-Oct-05    | 12                   | 126.6            | 26.70           | 0.22        | 0.16                  | 4.27                      | 55.8               | 1.4                   | 76.4                   |
|                |                   | 28-Sep-06    | 43 (44)              | 90.8             | 11.30           | 0.80 (0.82) | 0.55 (0.57)           | 6.22 (6.45)               | 53.7               | 1.5                   | 77.9                   |
|                | Venema Cr (upper) | 3-Mar-05     | 7                    | 104.1            | 15.40           | 0.14        | 0.10                  | 1.54                      | 49.0               | 1.5                   | 73.5                   |
|                |                   | 19-Oct-05    | 7                    | 106.4            | 14.50           | 0.14        | 0.07                  | 1.02                      | 49.9               | 1.9                   | 94.8                   |
|                |                   | 29-Sep-06    | 7                    | 79.4             | 6.00            | 0.14        | 0.12                  | 0.72                      | 51.6               | 1.1                   | 58.3                   |
| Thornton Creek | South Branch      | 24-Feb-05    | 241                  | 98.3             | 10.10           | 1.90        | 0.69                  | 6.97                      | 126.9              | 2.8                   | 351.5                  |
|                |                   | 13-Oct-05    | 309 (310)            | 87.6             | 8.34            | 2.70 (2.70) | 1.06 (1.06)           | 8.77 (8.77)               | 114.8              | 2.6                   | 293.9                  |
|                |                   | 2-Aug-06     | 489                  | 50.7             | 3.44            | 4.80        | 1.76                  | 6.05                      | 101.9              | 2.7                   | 277.2                  |
|                | Kramer Creek      | 25-Feb-05    | 76 (85)              | 108.0            | 15.80           | 0.70 (0.79) | 0.57 (0.64)           | 9.01 (9.67)               | 107.5              | 1.2                   | 133.3                  |
|                |                   | 28-Sep-05    | 31 (32)              | 121.4            | 24.80           | 0.29 (0.30) | 0.25 (0.26)           | 6.20 (6.25)               | 107.3              | 1.2                   | 123.4                  |
|                |                   | 26-Jul-06    | 52 (55)              | 87.2             | 11.15           | 0.50 (0.53) | 0.43 (0.45)           | 4.79 (5.45)               | 104.6              | 1.2                   | 121.3                  |




FIGURE 3.1.-- Length frequency (10-mm FL increments) of cutthroat trout collected from the reference site of Piper's Creek and lower Venema Creek. Total number of cutthroat trout sampled and mean length is given in each panel.



The two reference sites on Venema Creek were sampled three times; each time within a day of sampling Piper's Creek. The lower Venema site had a mean wetted width of 1.45 m and a maximum depth of 0.61 m. Stream habitat at this site was composed of approximately 32% pools (by area), 62% riffles and 1% glides. For the three dates combined, 85 fish were collected (does not include several chum salmon fry that originated from an acclimation pond), 84 were cutthroat trout and one was a rainbow trout. The estimated populations of cutthroat trout varied from 12 on October 19, 2005 to 43 on September 29, 2006. Overall, 59% of the cutthroat trout were collected in pools yet only 32% of the habitat by area was composed of pools.

The upstream reference site of Venema Creek was originally sampled in March of 2005 from Rm 200 to 250 but was later moved further upstream to Rm 270-320 for the October 2005 and September 2006 sampling periods. Only cutthroat trout were collected during each sampling period. The estimated population size ( $\hat{N} = 7$ ) was the same for each sampling period. For all surveys combined, 90.5% of the cutthroat trout were less than 120 mm ( $N = 21$ ; range, 71-152 mm FL). On average, pools made up roughly 9% of the habitat area, but 33% of the fish were collected in pools.

A reference site was also established on Mohlendorph Creek between Rm 100 and 161 and was surveyed on October 19, 2005. However, because no fish were captured and there was little streamflow and available habitat, we did not resurvey this site. The site had a mean wetted width of 0.93 m and a maximum depth of 0.20 m. Stream habitat at this site was composed of approximately 70% riffles (by area) and 30% glides.

*Thornton Creek watershed.*— The reference site on the South Branch of Thornton Creek (Maple Leaf Reach) was substantially larger (307 m<sup>2</sup>) than any other reference site (range, 77-192 m<sup>2</sup>). This site was located approximately 550 m from where the South Branch merges with the North Branch. Little LWD was present at this site (Table 3.4). Habitats at this site were composed of approximately 8% pools (by area), 59% riffles, 33% glides (Table 3.5). Large numbers of cutthroat trout were collected on each sampling date. Except of one threespine stickleback, they were the only species captured at this site. The estimated abundance of cutthroat trout increased in each sampling period

from February 2005 to August 2006 (Table 3.3). Most cutthroat trout were between 50 and 100 mm FL (range, 39-200 mm FL; Figure 3.2).

Our other reference site in the Thornton Creek watershed was Kramer Creek, a small tributary of the South Branch of Thornton Creek. The lowest 160 m of the stream was used as a reference site; however 52.5 m of this length were in seven culverts, which we could not sample and were not included in the analysis. The downstream end of the site was a convergence pool with the South Branch of Thornton Creek. The mean wetted width of the entire site was 1.18 m and had a maximum depth of 0.7 m. Stream habitat at this site was composed of 23% pools (by area) and 77% glides (Table 3.5). Cutthroat trout was the primary species captured ( $n = 167$ ), however three threespine stickleback, nine pumpkinseed, and one juvenile coho salmon were also captured. The highest abundance of cutthroat trout was on February 25, 2005 ( $\hat{N} = 76$ ) and the lowest was on September 26, 2006 ( $\hat{N} = 31$ ). Therefore, this tributary may be better habitat for trout in the winter (high-flow conditions) than during the late summer (low-flow conditions).

TABLE 3.4.-- Number and density (number of pieces /100 m) of large woody debris (LWD) at reference sites in Seattle's streams, 2005-2006. Each piece of LWD was at least 2 m long and 10 cm at the midpoint.

| Stream system | Site Location    | Date Sampled | Length Sampled (m) | Total # of LWD | LWD / 100 m |
|---------------|------------------|--------------|--------------------|----------------|-------------|
| Piper's       | Mainstem         | 3-Mar-05     | 52                 | 2              | 3.8         |
|               |                  | 19-Oct-05    | 51.5               | 5              | 9.7         |
|               | Lower Venema Cr. | 3-Mar-05     | 51                 | 8              | 15.7        |
|               |                  | 19-Oct-05    | 55.8               | 7              | 12.5        |
|               | Upper Venema Cr. | 3-Mar-05     | 49                 | 1              | 2.0         |
|               |                  | 20-Oct-05    | 49.9               | 6              | 12.0        |
|               |                  | Mohlendorph  | 19-Oct-05          | 60.6           | 0           |
| Thornton      | South Branch     | 24-Feb-05    | 126.9              | 1              | 0.8         |
|               |                  | 13-Oct-05    | 114.8              | 2              | 1.7         |
|               |                  | 2-Aug-06     | 101.9              | 3              | 2.9         |
|               | Kramer Creek     | 25-Feb-05    | 107.5              | 0              | 0.0         |
|               |                  | 28-Sep-05    | 107.3              | 0              | 0.0         |
|               |                  | 26-Jul-06    | 104.6              | 0              | 0.0         |
| Longfellow    | Mainstem         | 7-Aug-06     | 71.8               | 15             | 20.9        |
| Fauntleroy    | Mainstem         | 18-Oct-05    | 103.3              | 31             | 30.0        |
|               |                  | 8-Aug-06     | 102.9              | 51             | 49.6        |
| Taylor        | Mainstem         | 17-Oct-05    | 119.2              | 10             | 8.4         |
|               |                  | 3-Aug-06     | 99.7               | 17             | 17.1        |




FIGURE 3.2.-- Length frequency (10-mm FL increments) of cutthroat trout collected from the reference site of South Branch Thornton Creek (Maple Leaf Site) and Kramer Creek. Total number of cutthroat trout sampled and mean length is given in each panel.

TABLE 3.5.-- Habitat information and number of fish collected in different habitat types of reference sites in the Thornton Creek watershed. %Fish is the percent of fish for each reference site that was collected in each habitat type (not adjusted by area). CV = convergence pool.

| Site location            | Date Sampled | Habitat Type | Number of Habitats | Combined Lengths (m) | Mean Wetted Width (m) | Area (m <sup>2</sup> ) | Mean Depth (m) | Maximum Depth (m) | # of Fish Captured | % Fish |
|--------------------------|--------------|--------------|--------------------|----------------------|-----------------------|------------------------|----------------|-------------------|--------------------|--------|
| South Branch Thornton Cr | 24-Feb-05    | Riffle       | 5                  | 73.5                 | 2.9                   | 213.0                  | 0.15           | 0.38              | 75                 | 32     |
|                          |              | Pool         | 3                  | 25.9                 | 2.7                   | 70.0                   | 0.25           | 0.45              | 81                 | 34     |
|                          |              | Glide        | 2                  | 27.5                 | 2.4                   | 66.0                   | 0.20           | 0.35              | 82                 | 34     |
|                          | 13-Oct-05    | Riffle       | 5                  | 56.6                 | 2.6                   | 144.3                  | 0.11           | 0.27              | 88                 | 29     |
|                          |              | Pool         | 1                  | 2.2                  | 2.4                   | 5.2                    | 0.15           | 0.32              | 7                  | 2      |
|                          |              | Glide        | 5                  | 56.0                 | 2.6                   | 145.0                  | 0.15           | 0.31              | 211                | 69     |
|                          | 2-Aug-06     | Riffle       | 5                  | 40.6                 | 2.8                   | 113.7                  | 0.15           | 0.31              | 108                | 22     |
|                          |              | Glide        | 3                  | 30.1                 | 3.1                   | 93.3                   | 0.17           | 0.37              | 208                | 43     |
|                          |              | Run          | 3                  | 31.2                 | 2.3                   | 71.8                   | 0.10           | 0.21              | 167                | 35     |
| Kramer Cr                | 25-Feb-05    | Pool         | 2                  | 8.2                  | 1.8                   | 14.7                   | 0.20           | 0.40              | 19                 | 24     |
|                          |              | Glide        | 5                  | 95.2                 | 1.1                   | 104.8                  | 0.20           | 0.45              | 37                 | 47     |
|                          |              | CV           | 1                  | 4.1                  | 2.8                   | 11.5                   | 0.25           | 0.50              | 23                 | 29     |
|                          | 28-Sep-05    | Pool         | 2                  | 8.9                  | 1.7                   | 15.2                   | 0.15           | 0.38              | 7                  | 22     |
|                          |              | Glide        | 5                  | 94.2                 | 1.0                   | 94.2                   | 0.28           | 0.38              | 7                  | 22     |
|                          |              | CV           | 1                  | 4.2                  | 3.5                   | 14.7                   | 0.40           | 0.88              | 18                 | 56     |
|                          | 26-Jul-06    | Pool         | 2                  | 7.8                  | 1.7                   | 13.3                   | 0.18           | 0.35              | 24                 | 43     |
|                          |              | Glide        | 5                  | 92.2                 | 1.0                   | 90.4                   | 0.13           | 0.46              | 15                 | 27     |
|                          |              | CV           | 1                  | 4.6                  | 3.9                   | 17.9                   | 0.40           | 0.70              | 17                 | 30     |

*Longfellow Creek.*— The reference site in Longfellow Creek was only sampled once, August 2006. The reference site was 71 m with a mean wetted width of 2.7 m and a maximum depth of 0.56 m. Habitat at this site was comprised of 42% pools (by area), 7% riffles, and 51% glides (Table 3.6). The estimated fish abundance was 88 coho salmon, 80 threespine stickleback, and 7 prickly sculpin (Table 3.7). Coho salmon lengths ranged between 60 and 110 mm FL (mean FL, 81.8 mm) and threespine stickleback between 20 and 80 mm FL (mean FL, 38.6 mm; Figure 3.3). The total fish biomass estimate was generally low compared to other reference sites in similar-sized streams (Piper’s Creek and South Branch Thornton Creek; Figure 3.4). Many of the fish were collected in glides at the lower end of the site that had undercut banks. Twenty-three percent of the fish were collected in pools, 1% in riffles and 76% in glides.

*Fauntleroy Creek.*-- The reference site on Fauntleroy Creek was sampled twice, once in 2005 and again in 2006. The reference site was located at Rm 200-303 and was 103 m long with an average wetted width of 1.37 m. On both occasions, few fish were collected and only juvenile coho salmon were collected. The estimated population size of coho salmon varied from 14 in October 2005 to 32 in August 2006. A large percentage of juvenile coho salmon were less than 90 mm for October 2005 and less than 80 mm for August 2006 (Figure 3.5). Stream habitat was composed of 7% pools (by area) 79% riffles, and 14% glides. Sixty-five percent of juvenile coho salmon were taken from pools, 18% from riffles, and 17% from glides.

*Taylor Creek.*— The Taylor Creek reference site was the only site to have large numbers of both cutthroat trout and juvenile coho salmon present (Table 3.7; Figure 3.6). Taylor Creek has an impassable barrier downstream of our reference site. Therefore, the juvenile coho salmon were outplanted and the cutthroat trout were from a resident population that does not have any input from the lake. For both sample dates, cutthroat trout represented close to 75% of the fish biomass, but only 43 to 64% of the population size. Similar to reference sites on Longfellow Creek and Fauntleroy Creek, fish biomass was also low in Taylor Creek. Stream habitat consisted of 60% riffles, 11% pools, and 29% glides, while average fish catch consisted of 22% in riffles, 29% in pools, and 49% in glides.

TABLE 3.6.-- Habitat information and number of fish collected in different habitat types of reference sites in the Longfellow Creek, Fauntleroy Creek, and Taylor Creek. %Fish is the percent of fish for each reference site that was collected in each habitat type (not adjusted by area).

| Site location | Date Sampled | Habitat Types | Number of Habitats | Combined Lengths (m) | Mean Wetted Width (m) | Area (m <sup>2</sup> ) | Mean Depth (m) | Maximum Depth (m) | # of Fish Captured | % Fish |
|---------------|--------------|---------------|--------------------|----------------------|-----------------------|------------------------|----------------|-------------------|--------------------|--------|
| Longfellow Cr | 7-Aug-06     | Riffle        | 1                  | 7.1                  | 1.9                   | 13.5                   | 0.15           | 0.34              | 2                  | 1      |
|               |              | Pool          | 3                  | 21.0                 | 3.8                   | 79.8                   | 0.25           | 0.56              | 42                 | 29     |
|               |              | Glide         | 4                  | 43.0                 | 2.3                   | 98.9                   | 0.20           | 0.42              | 101                | 70     |
| Fauntleroy Cr | 18-Oct-05    | Riffle        | 7                  | 70.6                 | 1.4                   | 98.8                   | 0.05           | 0.10              | 7                  | 50     |
|               |              | Pool          | 6                  | 16.0                 | 1.3                   | 20.8                   | 0.20           | 0.50              | 5                  | 36     |
|               |              | Glide         | 5                  | 16.7                 | 1.5                   | 25.1                   | 0.10           | 0.20              | 2                  | 14     |
|               | 8-Aug-06     | Riffle        | 6                  | 93.0                 | 1.3                   | 120.9                  | 0.15           | 0.40              | 25                 | 81     |
|               |              | Glide         | 3                  | 9.9                  | 1.4                   | 14.3                   | 0.10           | 0.20              | 6                  | 19     |
|               |              |               |                    |                      |                       |                        |                |                   |                    |        |
| Taylor Cr     | 17-Oct-05    | Riffle        | 10                 | 75.2                 | 1.5                   | 114.3                  | 0.09           | 0.12              | 4                  | 11     |
|               |              | Pool          | 2                  | 6.8                  | 2.6                   | 17.8                   | 0.20           | 0.45              | 13                 | 35     |
|               |              | Glide         | 7                  | 37.2                 | 1.6                   | 58.8                   | 0.15           | 0.30              | 20                 | 54     |
|               | 3-Aug-06     | Riffle        | 9                  | 62.8                 | 1.8                   | 113.0                  | 0.10           | 0.15              | 44                 | 33     |
|               |              | Pool          | 4                  | 8.4                  | 2.7                   | 22.5                   | 0.20           | 0.48              | 31                 | 23     |
|               |              | Glide         | 5                  | 28.5                 | 1.8                   | 51.3                   | 0.15           | 0.21              | 59                 | 44     |

TABLE 3.7.-- Population and density estimates of reference sites in Longfellow Creek, Fauntleroy Creek, and Taylor Creek. Population estimates were calculated using depletion techniques of Carl and Strub (1978). Biomass (fish (g)/m<sup>2</sup>) is calculated by multiplying Fish/m<sup>2</sup> by the average weight (g). Species captured were coho salmon (COH), cutthroat trout (CUT), threespine stickleback (STB), and prickly sculpin (PKS).

| Stream           | Date sampled | Species | Estimated population | Mean length (mm) | Mean weight (g) | Fish / m | Fish / m <sup>2</sup> | Fish (g) / m <sup>2</sup> | Length of Unit (m) | Mean wetted width (m) | Area (m <sup>2</sup> ) |
|------------------|--------------|---------|----------------------|------------------|-----------------|----------|-----------------------|---------------------------|--------------------|-----------------------|------------------------|
| Longfellow Creek | 7-Aug-06     | COH     | 88                   | 81.7             | 7.20            | 1.24     | 0.44                  | 3.17                      | 71.0               | 2.7                   | 192.4                  |
|                  |              | STB     | 80                   | 38.6             | 0.70            | 0.10     | 0.40                  | 0.28                      |                    |                       |                        |
|                  |              | PKS     | 7                    | 128.0            | 27.70           | 1.13     | 0.04                  | 1.11                      |                    |                       |                        |
| Fauntleroy Creek | 18-Oct-05    | COH     | 14                   | 89               | 8.60            | 0.14     | 0.10                  | 0.86                      | 103.3              | 1.4                   | 144.6                  |
|                  | 8-Aug-06     | COH     | 32                   | 71.5             | 5.00            | 0.31     | 0.23                  | 1.15                      | 102.9              | 1.34                  | 137.9                  |
| Taylor Creek     | 17-Oct-05    | COH     | 21                   | 80.0             | 6.10            | 0.18     | 0.11                  | 0.67                      | 119.2              | 1.6                   | 190.7                  |
|                  |              | CUT     | 16                   | 133.0            | 31.38           | 0.13     | 0.08                  | 2.51                      |                    |                       |                        |
|                  | 3-Aug-06     | COH     | 48                   | 41.3             | 3.60            | 0.48     | 0.29                  | 1.04                      | 99.7               | 1.7                   | 168.5                  |
|                  |              | CUT     | 86                   | 63.9             | 5.10            | 0.86     | 0.51                  | 2.60                      |                    |                       |                        |




FIGURE 3.3.-- Length frequency (10-mm FL increments) of threespine stickleback (open bars) and juvenile coho salmon (shaded bars) collected from the reference site of Longfellow Creek, August 7, 2006. Total number of fish sampled and mean length is also given.




FIGURE 3.4.-- Salmonid biomass estimates ( $g/m^2$ ) for different seasons at eight reference sites, 2005-2006. Error bars represent the range of two observations. If no error bar is shown, it indicates it was only sampled once for a particular season. S.B. Thornton = South Branch of Thornton Creek; ND = no data.






FIGURE 3.5.-- Length frequency (10-mm FL increments) of juvenile coho salmon collected from the reference site of Fauntleroy Creek, October 18, 2005 (shaded bars) and August 6, 2006 (open bars). Total number of fish sampled and mean length is also given.




FIGURE 3.6.-- Length frequency (10-mm FL increments) of coho salmon and cutthroat trout collected from the reference site of Taylor Creek. Total number of fish sampled and mean length is also given.

## Discussion

Our reference-site surveys provide essential baseline information for future comparisons to examine changes after restoration activities or examine long-term changes in ecosystem health. Restoration work has been completed in the Maple Leaf Reach of the South Branch Thornton Creek and additional surveys are needed to determine the effectiveness of the restoration project. Our reference-site surveys in the Piper's Creek watershed are part of ongoing monitoring by NOAA Fisheries to assess a large natural drainage project. Restoration projects are in the planning stage for Kramer Creek (reconfigure stream channel) and Taylor Creek (remove barrier). In Longfellow Creek, ongoing studies are being conducted by NOAA Fisheries and USFWS to determine the cause of pre-mortality of adult coho salmon. If the cause is identified and stream habitat conditions improve, resampling of our reference site could provide valuable information.

The density of salmonids in our reference sites in the South Branch of Thornton Creek and Piper's Creek appears to be high in comparison to other lowland streams in the Pacific Northwest. For example, Roni (2000) sampled 30 lowland streams in Washington and Oregon during the summer low period using the same techniques we used. He used streams that had a bankfull width of 4 to 12 m. We did not measure bankfull width but our reference sites on South Branch of Thornton Creek (2.7 m wetted width) and Piper's Creek (2.8 wetted width) probably fit within this range. Using our summer samples for comparison, we observed a density of 4.8 salmonids/m in South Branch of Thornton Creek. Roni (2000) only observed a density higher than this in one stream (range, 0.22 to 5.03 salmonids/m). We did not sample Piper's Creek in the summer but we did sample in late September and we observed a density of 2.4 salmonids/m, which is higher than 80% of the streams sampled by Roni (2000). In the streams sampled by Roni (2000), coho salmon, steelhead, and cutthroat trout were present; whereas, in the South Branch of Thornton Creek and Piper's Creek almost all fish were cutthroat trout. The density of cutthroat trout (fish/m) in these two reference sites was approximately 1.2 to 96 times higher (assuming all unidentified juvenile trout were cutthroat trout) than in any stream sampled by Roni (2000).

Kramer Creek had an unexpectedly high abundance of fish. The creek is relatively small and has little woody debris or other structure. The creek runs along a road and is open

with no forested canopy (Figure 3.7). The stream is a series of plunge pools and glides separated by culverts that are under residential driveways. Many of the fish we collected in Kramer Creek were from the convergence pool with the South Branch of Thornton Creek. The convergence pool was relatively deep (maximum depth, 0.5-0.88 m), which may provide adequate cover for large cutthroat trout. Also, at the head end of each plunge pool, the depth was relatively deep, which may provide adequate trout habitat. The creek also had a large amount of water cress and other emergent vegetation, which may provide cover and food (macroinvertebrates) for trout (Figure 3.7). Research of streams in the Pacific Northwest has shown that open streams may be more productive and have a higher abundance of salmonids than streams with a forested canopy (Hawkins et al.1983).

Of the three sample dates, cutthroat trout abundance in Kramer Creek was highest during the winter sample. In contrast, winter cutthroat trout abundance in South Branch of Thornton Creek was slightly lower than the summer or fall samples. The abundance of other species (pumpkinseed, threespine stickleback, and juvenile coho salmon) in Kramer Creek was also highest in the winter. The lower part of Kramer Creek may provide good winter habitat because it has a low gradient and may not have high water velocities during storm events. Also, salmonids generally seek cover during the winter and the emergent vegetation in Kramer Creek may provide the necessary cover. Winter flow conditions in the South Branch of Thornton Creek may cause some cutthroat trout to move into smaller tributaries such as Kramer Creek.




FIGURE 3.7.-- Photo of the upstream end of the Kramer Creek reference site, July 2006. Water flows from the top of the photo to the bottom of the photo.

## CHAPTER 4. FISH INDEX OF BIOTIC INTEGRITY

### Introduction and Methods

To assess ecosystem health, we used a fish index of biotic integrity (FIBI) that has been developed for Puget Sound lowland streams (Matzen and Berge 2008). This particular FIBI was developed from 70 sites in 30 subbasins in the Lake Washington basin, which included three sites in Thornton Creek. The index has six metrics and each metric is scored from 1 (lowest) to 4 (highest) (Table 4.1). Therefore, FIBI scores can range from 6 to 24. The index was developed for second- and third-order streams and may not be useful for first-order streams because few fish species are usually present in first-order streams even under pristine conditions. We calculated FIBI scores for each index and reference site in second- and third-order streams where fish were present.

TABLE 4.1.— Metric scoring of the fish index of biotic integrity (FIBI) of Matzen and Berge (2008). Scores are determined by the percentage of the total number of fish sampled.

| Metrics  | Score |        |       |     |
|--|-------|--------|-------|-----|
|  | 1     | 2      | 3     | 4   |
| Percent invertivore individuals                  | <35   | 35-55  | 55-75 | ≥75 |
| Percent invertivore/piscivore individuals        | ≥65   | 45-65  | 25-45 | <25 |
| Percent coho salmon individuals                  | <5    | 5-25   | 25-41 | ≥41 |
| Percent cutthroat trout individuals              | ≥65   | 45-65  | 25-45 | <25 |
| Percent sculpin individuals                      | <0.5  | 0.5-10 | 10-40 | ≥40 |
| Percent individuals of the most abundant species | ≥80   | 65-80  | 50-65 | <50 |

## Results

Overall, FIBI scores were generally low, primarily due to the relatively high abundance of cutthroat trout and lack of other species such as coho salmon and cottids (Tables 4.2 and 4.3). Half of the FIBI scores for the index sites were less than 10. FIBI scores were generally higher in Longfellow Creek and Fauntleroy Creek than other Seattle streams, largely because few cutthroat trout were present and coho salmon were common. However, the FIBI scores from these streams may have been artificially high because juvenile coho salmon may have been outplanted. FIBI scores of Taylor Creek may have also been high due to outplantings of juvenile coho salmon. Within each basin, FIBI scores were usually highest close to the mouth of the creek and scores were often lowest at upstream locations. The presence of cottids and other fish species (besides cutthroat trout) close to the mouth of creek resulted in higher scores.

TABLE 4.2. – Fish index of biotic integrity (FIBI) scores and number of fish caught at reference sites in Seattle streams. FIBI scores can range from 6 to 24. Location (Rm) is the downstream end of the site. COH = juvenile coho salmon, CUT = cutthroat trout; SAL = other salmonids; STB = threespine stickleback; COT = cottids.

| Stream system # | Stream                    | Location (Rm) | Date sampled | FIBI score | Number caught |     |     |     |     |
|-----------------|---------------------------|---------------|--------------|------------|---------------|-----|-----|-----|-----|
|                 |                           |               |              |            | COH           | CUT | SAL | STB | COT |
| 5               | Piper's Cr.               | 800           | 3-Mar-05     | 6          |               | 27  |     |     |     |
|                 |                           |               | 19-Oct-05    | 9          | 1             | 43  |     |     |     |
|                 |                           |               | 29-Sep-06    | 6          |               | 112 |     |     |     |
| 5               | Lower Venema Cr.          | 0             | 3-Mar-05     | 6          |               | 29  |     |     |     |
|                 |                           |               | 19-Oct-05    | 6          |               | 12  |     |     |     |
|                 |                           |               | 29-Sep-06    | 6          |               | 43  | 1   |     |     |
| 14              | South Branch Thornton Cr. | 550           | 24-Feb-05    | 6          |               | 238 |     |     |     |
|                 |                           |               | 13-Oct-05    | 6          |               | 306 |     | 1   |     |
|                 |                           |               | 2-Aug-06     | 6          |               | 483 |     |     |     |
| 34              | Longfellow Cr.            | 100           | 7-Aug-06     | 20         | 89            | 1   |     | 59  | 7   |
| 43              | Fauntleroy Cr.            | 250           | 18-Oct-05    | 18         | 14            |     |     |     |     |
|                 |                           |               | 8-Aug-06     | 18         | 31            |     |     |     |     |
| 49              | Taylor Cr.                | 1,500         | 17-Oct-05    | 13         | 69            | 102 |     |     |     |
|                 |                           |               | 3-Aug-06     | 13         | 49            | 86  |     |     |     |

TABLE 4.3. – Fish index of biotic integrity (FIBI) scores and number of fish caught at index sites in Seattle streams, 2005. FIBI scores can range from 6 to 24. Location (Rm) is the downstream end of the site. LPU = lamprey, unidentified ammocoetes; COH = juvenile coho salmon, CUT = cutthroat trout; SAL = other salmonids; STB = threespine stickleback; CEN = centrarchids; COT = cottids.

| Stream system # | Stream                    | Location (Rm) | FIBI score | Number caught |     |     |     |     |     |     |
|-----------------|---------------------------|---------------|------------|---------------|-----|-----|-----|-----|-----|-----|
|                 |                           |               |            | LPU           | COH | CUT | SAL | STB | CEN | COT |
| 5               | Piper's Cr.               | 0             | 17         |               | 1   | 15  |     |     |     | 40  |
|                 |                           | 250           | 9          |               | 6   | 71  |     |     |     |     |
|                 |                           | 500           | 9          |               | 3   | 47  |     |     |     |     |
|                 |                           | 1,175         | 6          |               |     | 55  |     |     |     |     |
| 14              | Lower Thornton Cr.        | 0             | 21         |               | 6   |     |     | 3   |     | 74  |
|                 |                           | 250           | 15         |               |     | 7   |     |     |     | 120 |
|                 |                           | 350           | 18         | 4             | 1   | 25  |     |     | 1   | 312 |
|                 |                           | 500           | 13         |               | 9   | 135 |     | 43  | 5   | 7   |
|                 |                           | 1,100         | 14         |               | 1   | 67  |     | 31  | 17  | 1   |
|                 |                           | 1,900         | 12         |               | 7   | 421 |     |     | 5   | 1   |
| 14              | North Branch Thornton Cr. | 2,900         | 6          |               |     | 142 |     |     | 2   |     |
|                 |                           | 3,650         | 6          |               |     | 111 |     |     | 4   |     |
|                 |                           | 4,300         | 6          |               |     | 80  |     |     | 2   |     |
|                 |                           | 5,150         | 6          |               |     | 42  |     |     |     |     |
|                 |                           | 5,900         | 6          |               |     | 15  |     |     | 1   |     |
|                 |                           | 6,650         | 6          |               |     | 125 |     |     | 28  |     |
| 14              | South Branch Thornton Cr. | 0             | 6          |               |     | 51  |     |     |     |     |
|                 |                           | 1,350         | 6          |               |     | 90  |     |     |     |     |
|                 |                           | 2,250         | 7          |               |     | 29  |     | 13  |     |     |
|                 |                           | 3,200         | 7          |               |     | 17  | 1   | 4   |     |     |
|                 |                           | 3,750         | 12         |               |     | 1   |     | 78  |     |     |
| 34              | Longfellow Cr.            | 0             | 18         |               |     |     |     | 17  |     | 3   |
|                 |                           | 2,150         | 9          |               |     |     | 3   |     |     |     |
|                 |                           | 2,850         | 17         |               | 2   |     | 2   |     | 1   |     |
|                 |                           | 3,400         | 9          |               |     |     | 1   |     |     |     |
| 42              | Fautleroy Cr.             | 0             | 23         |               | 4   |     |     |     |     | 4   |
|                 |                           | 150           | 18         |               | 2   |     |     |     |     |     |
|                 |                           | 450           | 18         |               | 4   |     |     |     |     |     |
|                 |                           | 725           | 18         |               | 9   |     |     |     |     |     |
| 49              | Taylor Cr.                | 0             | 19         |               | 20  | 30  | 2   | 17  |     | 33  |
|                 |                           | 1,500         | 15         |               | 51  | 9   |     |     |     |     |
|                 |                           | 2,800         | 9          |               | 1   | 5   |     |     |     |     |

## Discussion

The FIBI of Matzen and Berge (2008) was developed to evaluate the relationship between urbanization and fish assemblages in Puget Sound lowland streams. As expected, FIBI scores of Seattle's streams were generally low and reflected the high degree of urbanization in each watershed. Matzen and Berge (2008) found a strong negative relationship between FIBI scores and percent total impervious area (TIA). For the five major watersheds, percent TIA ranges from 38 to 59% (City of Seattle 2007). Based on a regression developed by Matzen and Berge (2008), FIBI scores in these watersheds should range from approximately 6 to 12. Nineteen of 35 sites (54%) we sampled were close to the expected value (Figure 4.1).

However, at some sites, scores were higher than expected given the amount of urbanization (Figure 4.1). A regression developed from all of our sampling sites indicated a different relationship between FIBI scores and TIA than the relationship observed by Matzen and Berge (2008). The FIBI of Matzen and Berge (2008) may not work in some urban areas where the overall abundance of fish is generally low and many of the fish are planted. The FIBI scores are based on percentages and in streams with low fish abundance, slight differences in the catch could substantially change the FIBI score. Outplanting of juvenile coho salmon and the mysterious lack of cutthroat trout in Longfellow Creek and Fauntleroy Creek probably resulted in higher than expected FIBI scores. FIBI scores were also higher than expected near the mouth of each creek. These scores were often high because large numbers of prickly sculpin and coastrange sculpin were present that commonly move upstream from estuaries and lakes. Because these species are more tolerant of warm water temperatures than most other cottid species (Zaroban et al. 1999) and generally only inhabit the lower reaches of small streams, they may not be the best indicator of ecosystem health. The presence or absence of less tolerant cottid species that must complete their entire life cycle in the stream would probably be a better metric. Also, our results underscore the need to sample several sites in each basin and not just rely on scores from one site near the creek mouth.






FIGURE 4.1. Relationship between fish index of biotic integrity (FIBI scores, open circles,  $n = 35$ ) and percent total impervious area (%TIA) and for five basins with the City of Seattle. The lower regression line (solid line) is the predicted line based on sampling by Matzen and Berge (2008); the upper line is based on observed FIBI scores in City of Seattle streams. Values used for %TIA include: Fauntleroy Creek – 38%, Taylor Creek – 48%, Longfellow Creek – 52%, Piper’s Creek basin – 57% and Thornton Creek basin – 59%. Some %TIA values were altered slightly for display purposes (reduce overlapping values).

## **ACKNOWLEDGMENTS**

We wish to thank Joe Starstead, City of Seattle; Matthew Arasim, Steven Celedonia, Steve Damm, Ian Grettenberger, Tracy Leavy, Fan Lee, and Eric Tallman of the U.S. Fish and Wildlife Service (USFWS); and Sarah Morley, NOAA Fisheries for all their assistance with the field work. Katherine Lynch, Joe Starstead, Michele Koehler, and Keith Kurko, City of Seattle provided technical support and reviewed earlier drafts of this report. We also wish to thank the many landowners who allow us access through their property. Funding for this project was provided by the City of Seattle.

## REFERENCES

- Adams, S.B. and D.A. Schmetterling. 2007. Freshwater sculpins: phylogenetics to ecology. *Transactions of the American Fisheries Society* 136:1736-1741.
- Carle, F.L. and M.R. Strub. 1978. A new method for estimating population size from removal data. *Biometrics* 34:621-630.
- City of Seattle. 2007. State of the waters 2007, volume I: Seattle watercourses. Seattle, Washington.
- Erman, D. C., E. D. Andrews, and M. Yoder-Williams. 1988. Effects of winter floods on fishes in the Sierra Nevada. *Canadian Journal of Fisheries and Aquatic Sciences* 45:2195-2200.
- Foote, C. J. and G. S. Brown. 1998. Ecological relationship between freshwater sculpins (genus *Cottus*) and beach spawning sockeye salmon (*Oncorhynchus nerka*) in Iliamna Lake, Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* 55:1524-1533.
- Fransen, B.R., R.E. Bilby, S. Needham, and J.K. Walter. 1998. Delineating fish habitat based on physical characteristics associated with the upper extent of fish distributions. Paper presented at the 1998 Annual General Meeting, North Pacific International Chapter, American Fisheries Society, Alderbrook Inn, Union, Washington.
- Gray, M.A, R.A. Cunjak, and K.R. Munkittrick. 2004. Site fidelity of slimy sculpin (*Cottus cognatus*): insights from stable carbon and nitrogen analysis. *Canadian Journal of Fisheries and Aquatic Sciences* 61:1717-1722.
- Gray, M.A, R.A. Curry, and K.R. Munkittrick. 2005. Impacts of nonpoint inputs from potato farming on the populations of slimy sculpin (*Cottus cognatus*). *Environmental Toxicology and Chemistry* 24:2291-2298.
- Hawkins, C.P., M.L. Murphy, N.H. Anderson, and M.A. Wilzbach. 1983. Density of fish and salamanders in relation to riparian canopy and physical habitat in streams of the Northwestern United States. *Canadian Journal of Fisheries and Aquatic Sciences* 1173-1185.
- Horner, R. R. and C. W. May. 1998. Watershed urbanization and the decline of salmon in Puget Sound streams. Pages 19-40 *in* Salmon in the City Conference, Mount Vernon , Washington, May 20-21, 1998.
- Karr, J.R. 1998. Rivers as sentinels: using the biology of rivers to guide landscape management. Pages 502-528 in R.J. Naiman and R.E. Bilby, editors. *River*



- ecology and management: lessons from the Pacific Coastal ecoregion. Springer, New York.
- Latterell, J.J, R.J. Naiman, B.R. Fransen, and P.A. Bisson. 2003. Physical constraints on trout (*Oncorhynchus* spp.) distribution in the Cascade Mountains: a comparison of logged and unlogged streams. *Canadian Journal of Fisheries and Aquatic Sciences* 60:1007-1017.
- Leavy, T.R., P. Bakke., and R.J. Peters. 2008. Influence of habitat complexity and floodplain reconnection projects on the physical and biological conditions of Seattle urban streams: pre-project monitoring, final progress report, 2005-2007. Report of the U.S. Fish and Wildlife Service, Western Washington Fish and Wildlife Office, Lacey, Washington.
- LeMoine, M. 2007. Barriers to upstream migration of prickly sculpin *Cottus asper* and coastrange sculpin *Cottus aleuticus*. Master's thesis, Western Washington University, Bellingham.
- Lowry, G.R. 1966. Production and food of cutthroat trout in three Oregon coastal streams. *Journal of Wildlife Management* 30:754-767.
- Ludwa, K., G. Lucchetti, K.L. Fresh, and K. Walter. 1997. Assessing stream-dwelling fishes in basins of the Lake Washington watershed, summer 1996. King County Department of Natural Resources, Seattle, Washington.
- Mason, J. C. and S. Machodori. 1976. Populations of sympatric sculpins, *Cottus aleuticus* and *Cottus asper*, in four adjacent salmon-producing coastal streams on Vancouver Island, B.C. *Fishery Bulletin* 74:131-141.
- Matzen, D.A. and H.B. Berge. 2008. Assessing small-stream biotic integrity using fish assemblages across an urban landscape in the Puget Sound lowlands of western Washington. *Transactions of the American Fisheries Society* 137:677-689.
- McPhail, J.D. 2007. The freshwater fishes of British Columbia. University of Alberta Press, Edmonton.
- MNDNR (Minnesota Department of Natural Resources). 2003. Southeast Minnesota sculpin reintroduction plan. Available: [files.dnr.state.us/areas/fisheries/lakecity/semn\\_sculpin\\_reintrplan.pdf](http://files.dnr.state.us/areas/fisheries/lakecity/semn_sculpin_reintrplan.pdf) (May 2007).
- Morrow, J. E. 1980. The freshwater fishes of Alaska. Alaska Northwest Publishing Company, Anchorage.
- Moyle, P. B. 1977. In defense of sculpins. *Fisheries* 2(1):20-23.

- Moyle, P. B. 2002. Inland fishes of California. University of California Press, Berkeley.
- Mueller, K. W. 1998. 1997 Green Lake survey: the warmwater fish community of an urban lake plagued by algal blooms and Eurasian watermilfoil. Warmwater Enhancement Program, Washington Department of Fish and Wildlife, Olympia.
- Mueller, K. W. and M. R. Downen. 2000. 1999 Green Lake surveys: aspects of the biology of common carp with notes on the warmwater fish community. Warmwater Enhancement Program, Washington Department of Fish and Wildlife, Olympia.
- Muto, M. and L. Shefler. 1983. Game fish distribution in selected streams within the Lake Washington drainage basin. Report of Washington Department of Game to METRO, Water Quality Division, Seattle, Washington.
- Pfeifer, B. 1984. A survey of Piper's Creek in Seattle, Washington. Fisheries Management Division, Washington State Game Department, Olympia.
- Pleus, A.E. 1999. TFW Monitoring Program method manual for the wadeable stream discharge method. Report TFW-AM9-99-009 of Northwest Indian Fisheries Commission to Washington Department of Natural Resources, Olympia, Washington.
- Pleus, A.E. 1999. TFW Monitoring Program method manual for the habitat unit survey. Report TFW-AM9-99-003 of Northwest Indian Fisheries Commission to Washington Department of Natural Resources, Olympia, Washington.
- Price, B.E. 2006. The summer feeding ecology of rainbow trout (*Oncorhynchus mykiss*) and cutthroat trout (*O. clarkii*) in the Cedar River, Washington. Master's thesis, The Evergreen State College, Olympia, Washington.
- Rantz, S.E. and others. 1982. Measurement and computation of streamflow: volume I. Measurement of stage and discharge. U.S. Geological Survey Water-Supply Paper 2175. U.S. Government Printing Office, Washington, D.C.
- Reed, L., K. Lynch, K. Paulsen, B. McMillan, M. Myers, G. Ylitalo, J. Labenia, T. Collier, and N. Scholz. 2004. High rates of pre-spawn mortality in coho salmon (*Oncorhynchus kisutch*) in urban streams in Puget Sound basin. Proceedings of the 2003 Georgia Basin/Puget Sound Research Conference, Vancouver, British Columbia.
- Resource Planning Associates. 1998. Lower Thornton Creek evaluation of salmonid habitat conditions and opportunities. Task 1 memorandum, Report of Resource Planning Associates to Seattle Public Utilities, Seattle, Washington.

- Rosenfeld, J., M. Porter, and E. Parkinson. 2000. Habitat factors affecting the abundance and distribution of juvenile cutthroat trout (*Oncorhynchus clarki*) and coho salmon (*Oncorhynchus kisutch*). *Canadian Journal of Fisheries and Aquatic Sciences* 57:766-774.
- Schuett-Hames, D., A.E. Pleus, J. Ward, M. Fox, and J. Light. 1999. TFW Monitoring Program method manual for the large woody debris survey. Report TFW-AM9-99-004 of Northwest Indian Fisheries Commission to Washington Department of Natural Resources, Olympia, Washington.
- Scott, J.B., C. R. Steward, and Q. J. Stober. 1986. Effects of urban development on fish population dynamics in Kelsey Creek, Washington. *Transactions of the American Fisheries Society* 115:555-567.
- Seiler, D., L. Peterson, J. Serl, and R. Tabor. 2005. Cutthroat trout as successful urbanites. Proceeding of the 2005 coastal cutthroat trout symposium, status, management, biology, and conservation. Port Townsend, Washington, September 29-October 1, 2005.
- Serl, J. D. 1999. The response of stream fish communities of the Lake Washington watershed to various magnitudes of urbanization. Master's thesis, University of Washington, Seattle.
- Shapovalov, L. and A. C. Taft. 1954. The life histories of steelhead rainbow trout (*Salmo gairdneri gairdneri*) and silver salmon (*Oncorhynchus kitsch*). California Department of Fish and Game, Fish Bulletin 98.
- Tabor, R. A., K. L. Fresh, D. K. Paige, E. J. Warner, and R. J. Peters. 2007a. Distribution and habitat use of cottids in the Lake Washington basin. *American Fisheries Society Symposium* 53:25-40.
- Tabor, R.A., J. A. Schuerer, H. A. Gearns, and E. P. Bixler. 2004. Nearshore habitat use by juvenile Chinook salmon in lentic systems of the Lake Washington basin, annual report, 2002. U.S. Fish and Wildlife Service, Western Washington Fish and Wildlife Office, Lacey, Washington.
- Tabor, R.A., E. Warner, and S. Hager. 2001. An oriental weatherfish (*Misgurnus anguillicaudatus*) population established in Washington State. *Northwest Science* 75:72-76.
- Tabor, R.A., E. J. Warner, K.L. Fresh, B.A. Footen and J.R. Chan. 2007b. Ontogenetic diet shifts of prickly sculpin in the Lake Washington basin, Washington. *Transactions of the American Fisheries Society* 136:1801-1813.
- Thomas, G.L. 1992. Restoration of fish habitat in Piper's Creek, Seattle, Washington. Report of Prince William Sound Science Center to Municipality of Metropolitan



Seattle, Washington.

- Trotter, P.C. 2000. Headwater fishes and their uppermost habitats: a review as background for stream typing. Timber, Fish, and Wildlife Report TFW-ISAG1-00-001 to Washington Department of Natural Resources, Olympia, Washington.
- Utzinger, J., C. Roth, and A. Peter. Effects of environmental parameters on the distribution of bullhead *Cottus gobio* with particular consideration of the effects of obstructions. *Journal of Applied Ecology* 35:882-892.
- Washington Trout. 2000. Water typing and fish distribution within the City of Seattle. Draft report, May 18, 2000.
- Welsh, H.H. Jr. and L.M. Ollivier. 1998. Stream amphibians as indicators of ecosystem stress: a case study from California's redwoods. *Ecological Applications* 8:1118-1132.
- Wydoski, R.S., and R.R. Whitney. 2003. Inland fishes of Washington. University of Washington Press, Seattle.
- Zaroban, D.W., M.P. Mulvey, T.R. Maret, R.M. Hughes, and G.D. Merrit. 1999. Classification of species attributes for Pacific Northwest freshwater fishes. *Northwest Science* 73:81-92.
- Zimmerman, J.K.H. and B. Vondracek. 2007. Interactions between slimy sculpin and trout: slimy sculpin growth and diet in relation to native and nonnative trout. *Transactions of the American Fisheries Society* 136:1971-1800.





APPENDIX 1.1.-- Stream typing results and sample locations (fish distribution and habitat surveys) for stream system #'s 1-4. All sampling occurred in October 2005. Symbols of sample locations indicate the downstream end of each study reach.








APPENDIX 1.2.-- Stream typing results and sample locations for lower Piper's Creek (stream system #5). Sample location numbers from 1 to 299 are summer fish distribution surveys, 300 to 350 are winter fish distribution surveys, and 500 to 560 are reference site sampling. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.3.-- Stream typing results and sample locations for upper Piper's Creek (stream system #5). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.4.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Blue Ridge Creek (stream system #8) and stream system #9. All sampling occurred in October 2005. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.5.-- Stream typing results and sample locations (fish distribution and habitat surveys) for stream system #'s 10-11. All sampling occurred in October 2005. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.6.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Golden Garden streams (stream system #12). All sampling occurred in October 2005. Symbols of sample locations indicate the downstream end of each study reach.



APPENDIX 1.7.-- Stream typing results and sample locations for Licton Springs Creek (stream system #13). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.8.-- Stream typing results and sample locations for lower Thornton Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.9.-- Stream typing results and sample locations for lower Thornton Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.








APPENDIX 1.10.-- Stream typing results and sample locations for lower Thornton Creek and the lower South Branch of Thornton Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys, 300 to 350 are winter fish distribution surveys, and 500 to 560 are reference site sampling. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.11.-- Stream typing results and sample locations for the lower North Branch of Thornton Creek and Littlebrook Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.12.-- Stream typing results and sample locations for the North Branch of Thornton Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.13.-- Stream typing results and sample locations for the upper North Branch of Thornton Creek (including the Jackson Park Golf Course ponds) and Littles Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.14.-- Stream typing results and sample locations for the lower South Branch of Thornton Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 500 to 560 are reference site sampling. Symbols of sample locations indicate the downstream end of each study reach.



APPENDIX 1.15.-- Stream typing results and sample locations for Willow Creek and other tributaries to the lower South Branch of Thornton Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.16.-- Stream typing results and sample locations for upper South Branch of Thornton Creek (stream system #14). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.17.-- Stream typing results for Yesler Creek (stream system # 16).







APPENDIX 1.18.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Ravenna Creek (stream system # 17) and Iverness Creek (stream system # 15). All sampling occurred in September-October 2005. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.19.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Lawton Creek (stream system # 19), Mahteen Creek (stream system # 18), and Wolfe Creek (stream system # 20). All sampling occurred in August 2005. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.20.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Scheuerman Creek (stream system # 21), Owl's Creek (stream system # 22), and stream systems #'s 23-26. All sampling occurred in August-October 2005. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.21.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Interlaken streams (stream system #'s 28-30) and Washington Park stream (stream system # 27). All sampling occurred in August 2005. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.22.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Madrona Park streams (stream system #'s 31-32) and Frink Creek (stream system # 33). All sampling occurred in July-August 2005. Symbols of sample locations indicate the downstream end of each study reach.



APPENDIX 1.23.-- Stream typing results and sample locations for lower Longfellow Creek (stream system #34). Sample location numbers from 1 to 299 are summer fish distribution surveys, 300 to 350 are winter fish distribution surveys, and 500 to 560 are reference site sampling. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.24.-- Stream typing results and sample locations for upper Longfellow Creek (stream system #34). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.





APPENDIX 1.25.-- Stream typing results and sample locations for Puget Creek (stream system #35). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.







APPENDIX 1.26.-- Stream typing results and sample locations (fish distribution and habitat surveys) for Schmitz Creek (stream system # 38), Fairmount Creek (stream system # 37), and stream system # 36. All sampling occurred in August 2005. Symbols of sample locations indicate the downstream end of each study reach.




APPENDIX 1.27.-- Stream typing results and sample locations for Fautleroy Creek (stream system #42), Pelly Creek (stream system #40), and Mee-Kwa-Mooks Creek (stream system #39). Sample location numbers from 1 to 299 are summer fish distribution surveys and 500 to 560 are reference site sampling. Symbols of sample locations indicate the downstream end of each study reach.



APPENDIX 1.28.-- Stream typing results and sample locations for Seola Beach Creek (stream system # 43) and Durham Creek streams (stream system #'s 44-46). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.



APPENDIX 1.29.-- Stream typing results and sample locations for Mapes Creek (stream system # 48) and Mt Baker Creek (stream system # 47). Sample location numbers from 1 to 299 are summer fish distribution surveys and 300 to 350 are winter fish distribution surveys. Symbols of sample locations indicate the downstream end of each study reach.



APPENDIX 1.30.-- Stream typing results and sample locations for Taylor Creek (stream system #49). Sample location numbers from 1 to 299 are summer fish distribution surveys, 300 to 350 are winter fish distribution surveys, and 500 to 560 are reference site sampling. Symbols of sample locations indicate the downstream end of each study reach.